

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
13 November 2003 (13.11.2003)

PCT

(10) International Publication Number  
**WO 03/092588 A2**

(51) International Patent Classification<sup>7</sup>: **A61K**

(21) International Application Number: PCT/US03/13120

(22) International Filing Date: 25 April 2003 (25.04.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/376,779 30 April 2002 (30.04.2002) US

(71) Applicant (*for all designated States except US*): **MERCK & CO., INC.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **WANG, Ji-abing** [CN/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **MCVEAN, Carol, A.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(74) Common Representative: **MERCK & CO., INC.**; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SI, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— *without international search report and to be republished upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: 4-AZASTEROID DERIVATIVES AS ANDROGEN RECEPTOR MODULATORS

(57) Abstract: Compounds of structural formula I are modulators of the androgen receptor (AR) in a tissue selective manner. They are useful as agonists of the androgen receptor in bone and/or muscle tissue while antagonizing the AR in the prostate of a male patient or in the uterus of a female patient. These compounds are therefore useful in the treatment of conditions caused by androgen deficiency or which can be ameliorated by androgen administration, including osteoporosis, osteopenia, glucocorticoid-induced osteoporosis, periodontal disease, bone fracture, bone damage following bone reconstructive surgery, sarcopenia, frailty, aging skin, male hypogonadism, postmenopausal symptoms in women, atherosclerosis, hypercholesterolemia, hyperlipidemia, obesity, aplastic anemia and other hematopoietic disorders, inflammatory arthritis and joint repair, HIV-wasting, prostate cancer, cancer cachexia, Alzheimer's disease, muscular dystrophies, premature ovarian failure, and autoimmune disease, alone or in combination with other active agents.

WO 03/092588 A2

## TITLE OF THE INVENTION

## 4-AZASTEROID DERIVATIVES AS ANDROGEN RECEPTOR MODULATORS

## FIELD OF THE INVENTION

5                   The present invention relates to 4-azasteroid derivatives, their synthesis, and their use as androgen receptor modulators. More particularly, the compounds of the present invention are tissue-selective androgen receptor modulators and are thereby useful for the treatment of conditions caused by androgen deficiency or which can be ameliorated by androgen administration, such as osteoporosis,  
10   periodontal disease, bone fracture, frailty, and sarcopenia.

## BACKGROUND OF THE INVENTION

                  The androgen receptor (AR) belongs to the superfamily of steroid/thyroid hormone nuclear receptors, whose other members include the estrogen  
15   receptor (ER), the progesterone receptor (PR), the glucocorticoid receptor (GR), and the mineralocorticoid receptor (MR). The AR is expressed in numerous tissues of the body and is the receptor through which the physiological as well as the pathophysiological effects of endogenous androgen ligands, such as testosterone (T) and dihydrotestosterone (DHT), are expressed. Structurally, the AR is composed of  
20   three main functional domains: the ligand binding domain (LBD), the DNA-binding domain, and amino-terminal domain. A compound that binds to the AR and mimics the effects of an endogenous AR ligand is referred to as an AR agonist, whereas a compound that inhibits the effects of an endogenous AR ligand is termed an AR antagonist.

25                   Androgen ligand binding to the AR affords a ligand/receptor complex, which, subsequent to translocation inside the nucleus of the cell, binds to specific regulatory DNA sequences (referred to as androgen response elements or AREs) within the promoter or enhancer regions of the target gene or genes present in the cell's nucleus. Other proteins termed cofactors are next recruited which bind to the

amino-terminal domain or the LBD of the receptor leading to gene transcription and subsequent translation to produce the protein(s) encoded by that gene or genes.

Androgen therapy has been used in the clinic to treat a variety of male disorders, such as reproductive disorders and primary or secondary male hypogonadism. Moreover, a number of natural or synthetic AR agonists have been clinically investigated for the treatment of musculoskeletal disorders, such as bone disease, hematopoietic disorders, neuromuscular disease, rheumatological disease, wasting disease, and for hormone replacement therapy (HRT), such as female androgen deficiency. In addition, AR antagonists, such as flutamide and bicalutamide, have been used to treat prostate cancer. It would therefore be useful to have available compounds that can activate ("agonize") the function of the AR in a tissue-selective manner which would afford the desired beneficial osteo- and myoanabolic effects of androgens but without the negative androgenic properties, such as virilization and induction of an atherogenic lipid profile which can lead to cardiovascular disease.

The role of androgens in bone formation has been documented. For example, anabolic steroids, such as nandrolone decanoate or stanozolol, have been shown to increase bone mass in postmenopausal women. The beneficial effects of androgens on bone in postmenopausal osteoporosis were documented in recent studies using combined testosterone and estrogen administration [Hofbauer, et al., "Androgen effects on bone metabolism: recent progress and controversies," Eur. J. Endocrinol. 140: 271-286 (1999)]. Combined treatment significantly increased the rate and extent of the rise in bone mineral density (BMD) in the lumbar and hip regions, relative to treatment with estrogen alone. Additionally, estrogen - progestin combinations that incorporated an androgenic progestin (such as norethindrone), rather than medroxyprogesterone acetate, yielded greater improvements in hip BMD. These results have recently been confirmed in a larger 2-year, double-blind comparison study in which oral conjugated estrogen (CEE) and methyltestosterone combinations were demonstrated to be effective in promoting accrual of bone mass in the spine and hip, while conjugated estrogen therapy alone prevented bone loss ["A

two-year, double-blind comparison of estrogen-androgen and conjugated estrogens in surgically menopausal women: Effects on bone mineral density, symptoms and lipid profiles,” J. Reprod. Med., 44: 1012-1020 (1999)]. Despite the beneficial effects of androgens in postmenopausal women, the use of androgens has been limited because  
5 of the undesirable virilizing and metabolic action of androgens. The data from Watts and colleagues demonstrate that hot flushes decrease in women treated with CEE and methyltestosterone; however, 30% of these women suffered from significant increases in acne and facial hair, a complication of all current androgen pharmacotherapies [Watts, et al., “Comparison of oral estrogens and estrogens plus androgen on bone  
10 mineral density, menopausal symptoms, and lipid-lipoprotein profiles in surgical menopause,” Obstet. Gynecol., 85: 529-537 (1995)]. Moreover, the addition of methyltestosterone to CEE markedly decreased HDL levels, as seen in other studies. Therefore, non-tissue selective AR agonists may increase the risk of cardiovascular disease. Thus, the virilizing potential and negative effects on lipid profile of current  
15 androgen therapies provide a strong rationale for developing tissue-selective androgen receptor agonists for bone. Reference is made to J. A. Kanis, “Other agents for generalized osteoporosis,” in Osteoporosis, Blackwell Science, Ch. 8, pp 196-227 (1994) for a discussion of non-selective anabolic steroids in the treatment of osteoporosis.

20               It is also well established that androgens play an important role in bone metabolism in men, which parallels the role of estrogens in women [Anderson, et al., “Androgen supplementation in eugonadal men with osteoporosis – effects of six months of treatment on bone mineral density and cardiovascular risk factors,” Bone, 18: 171-177 (1996)]. Even in eugonadal men with established osteoporosis, the  
25 therapeutic response to testosterone treatment provided additional evidence that androgens exert important osteoanabolic effects. Mean lumbar BMD increased from 0.799 gm/cm<sup>2</sup> to 0.839 g/cm<sup>2</sup>, in 5 to 6 months in response to 250 mg of testosterone ester administered intramuscularly every fortnight. A common scenario for androgen deficiency occurs in men with stage D prostate cancer (metastatic) who undergo  
30 androgen deprivation therapy (ADT). Endocrine orchiectomy is achieved by long

acting GnRH agonists, while androgen receptor blockade is implemented with flutamide, nilutamide, bicalutamide, or RU 58841 (AR antagonists). In response to hormonal deprivation, these men suffered from hot flushes, significant bone loss, weakness, and fatigue. In a recent pilot study of men with stage D prostate cancer, osteopenia (50% vs. 38%) and osteoporosis (38% vs. 25%) were more common in men who had undergone ADT for greater than one year than the patients who did not undergo ADT [Wei, et al., "Androgen deprivation therapy for prostate cancer results in significant loss of bone density," Urology, 54: 607-611 (1999)]. Lumbar spine BMD was significantly lower in men who had undergone ADT. Thus, in addition to the use of tissue selective AR agonists for osteoporosis, tissue selective AR antagonists in the prostate that lack antagonistic action in bone and muscle may be useful agents for the treatment of prostate cancer, either alone or as an adjunct to traditional ADT such as with a GnRH agonist/antagonist [See also A. Stoch, et al., J. Clin. Endocrin. Metab., 86: 2787-2791 (2001)]. Tissue-selective AR antagonists may also have utility in the treatment of polycystic ovarian syndrome in postmenopausal women [see C.A. Eagleson, et al., "Polycystic ovarian syndrome: evidence that flutamide restores sensitivity of the gonadotropin-releasing hormone pulse generator to inhibition by estradiol and progesterone," J. Clin. Endocrinol. Metab., 85: 4047-4052 (2000) and E. Diamanti-Kandarakis, "The Effect of a Pure Antiandrogen Receptor Blocker, Flutamide, on the Lipid Profile in the Polycystic Ovary Syndrome," Int. J. Endocrinol. Metab., 83: 2699-2705 (1998)].

There is a need for more effective agents to treat osteopenia and osteoporosis in both men and women. Osteoporosis is characterized by bone loss, resulting from an imbalance between bone resorption (destruction) and bone formation, which starts in the fourth decade and continues throughout life at the rate of about 1-4% per year [Eastell, "Treatment of postmenopausal osteoporosis," New Engl. J. Med., 338: 736 (1998)]. In the United States, there are currently about 20 million people with detectable fractures of the vertebrae due to osteoporosis. In addition, there are about 250,000 hip fractures per year due to osteoporosis, associated with a 12%-20% mortality rate within the first two years, while 30% of patients

require nursing home care after the fracture and many never become fully ambulatory again. In postmenopausal women, estrogen deficiency leads to increased bone resorption resulting in bone loss in the vertebrae of around 5% per year, immediately following menopause. Thus, first line treatment/prevention of this condition is

5 inhibition of bone resorption by bisphosphonates, estrogens, selective estrogen receptor modulators (SERMs), and calcitonin. However, inhibitors of bone resorption are not sufficient to restore bone mass for patients who have already lost a significant amount of bone. The increase in spinal BMD attained by bisphosphonate treatment can reach 11% after 7 years of treatment with alendronate. In addition, as the rate of

10 bone turnover differs from site to site, higher in the trabecular bone of the vertebrae than in the cortex of the long bones, the bone resorption inhibitors are less effective in increasing hip BMD and preventing hip fracture. Therefore, osteoanabolic agents, which increase cortical bone formation and bone mass of long bones by stimulating periosteal bone formation, would address an unmet need in the treatment of

15 osteoporosis especially for patients with high risk of hip fractures. The osteoanabolic agents also complement the bone resorption inhibitors that target the trabecular envelope, leading to a biomechanically favorable bone structure (Schmidt, et al., "Anabolic steroid: Steroid effects on bone in women," In: J. P. Bilezikian, et al., Ed., Principles of Bone Biology, San Diego: Academic Press, 1996). Tissue-selective AR

20 agonists with diminished deleterious effects on the cardiovascular system and limited virilizing potential may be useful as a monotherapy for the prevention and/or treatment of female osteoporosis. In addition, a compound with osteoanabolic properties in bone and muscle but with reduced activity in the prostate and sex accessory tissues may be useful for the prevention and/or treatment of male

25 osteoporosis and osteopenia in men, particularly elderly men.

Selective androgen receptor modulators may also be useful to treat certain hematopoietic disorders. It is known that androgens stimulate renal hypertrophy and erythropoietin (EPO) production. Prior to the introduction of recombinant human EPO, androgens were employed to treat anemia caused by

30 chronic renal failure. In addition, androgens at pharmacological doses were found to

increase serum EPO levels in anemic patients with non-severe aplastic anemia and myelodysplastic syndromes but not in non-anemic patients. Treatment modalities for anemia will require selective action such as may be provided by selective androgen receptor modulators.

5                   Furthermore, selective androgen receptor modulators may also have clinical value as an adjunct to the treatment of obesity. This approach to lowering body fat is supported by published observations that androgen administration reduced subcutaneous and visceral abdominal fat in obese men [J.C. Lovejoy, et al., "Oral anabolic steroid treatment, but not parenteral androgen treatment, decreases  
10   abdominal fat in obese, older men," Int. J. Obesity, 19: 614-624 (1995)]. Therefore, SARMs devoid of androgenic effects on prostate may be beneficial in the treatment of obese men. In a separate study, androgen administration resulted in loss of subcutaneous abdominal fat in obese postmenopausal women [J.C. Lovejoy, et al., "Exogenous Androgens Influence Body Composition and Regional Body Fat  
15   Distribution in Obese Postmenopausal Women – A Clinical Research Center Study," J. Clin. Endocrinol. Metab., 81: 2198-2203 (1996)]. In the latter study, nandrolone decanoate, a weak androgen and anabolic agent, was found to increase lean body mass and resting metabolic rate in obese postmenopausal women consuming a weight-reducing diet.

20                   That androgen receptor agonists may also have therapeutic value against neurodegenerative diseases such as Alzheimer's disease (AD) has also been suggested in the art. The ability of androgens to induce neuroprotection through the androgen receptor was reported by J. Hammond, et al., "Testosterone-mediated neuroprotection through the androgen receptor in human primary neurons," J.  
25   Neurochem., 77: 1319-1326 (2001). Gouras et al. have observed that testosterone can reduce neuronal secretion of Alzheimer's  $\beta$ -amyloid peptides and may therefore be protective in the treatment of AD [Proc. Nat. Acad. Sci., 97: 1202-1205 (2000)]. A mechanism via inhibition of hyperphosphorylation of proteins implicated in the progression AD has also been described [S. Papasozomenos, "Testosterone prevents  
30   the heat shock-induced overactivation of glycogen synthase kinase-3 $\beta$  but not of

cyclin-dependent kinase 5 and c-Jun NH<sub>2</sub>-terminal kinase and concomitantly abolishes hyperphosphorylation of  $\tau$ : Implications for Alzheimer's disease," Proc. Nat. Acad. Sci., 99: 1140-1145 (2002)].

Non-steroidal compounds having androgen receptor modulating properties were disclosed in U.S. Patent Nos. 5,688,808; 5,696,130; 6,017,924; 6,093,821; WO 01/16139 (published 8 March 2001); and WO 01/16108 (published 8 March 2001), all assigned to Ligand Pharmaceuticals, and in WO 01/27086, assigned to Kaken Pharm. Co. Additional background for the rationale behind the development of Selective Androgen Receptor Modulators is found in L. Zhi and E. Martinborough in Ann. Rep. Med. Chem. 36: 169-180 (2001). Non-steroidal SARMs were disclosed in J.P. Edwards, "New Nonsteroidal Androgen Receptor Modulators Based on 4-(Trifluoromethyl)-2(1H)-Pyrrolidino[3,2-g]quinolinone," Bioorg. Med. Chem. Lett., 8: 745-750 (1998) and in L. Zhi et al., "Switching Androgen Receptor Antagonists to Agonists by Modifying C-ring Substituents on Piperidino[3,4-g]quinolinone," Bioorg. Med. Chem. Lett., 9: 1009-1012 (1999).

There exists a need in the clinical art for more effective agents that can elicit the positive responses of androgen replacement therapy but without the undesired side effects of non-tissue selective agonists of the AR. What is needed are compounds that can produce the same positive responses as androgen replacement therapy but without the undesired side effects. Also needed are androgenic compounds that exert selective effects on different tissues of the body. In this invention, we have identified compounds that function as selective androgen receptor modulators (SARMs) using a series of *in vitro* cell-assays that profile ligand mediated activation of AR, such as (i) N-C interaction, (ii) transcriptional repression, and (iii) transcriptional activation. SARM compounds in this invention, identified with the methods listed above, exhibit tissue selective AR agonism *in vivo*, i.e. agonism in bone (stimulation of bone formation in a rodent model of osteoporosis) and antagonism in prostate (minimal effects on prostate growth in castrated rodents and antagonism of prostate growth induced by AR agonists).

The compounds of the present invention identified as SARMs are useful to treat diseases or conditions caused by androgen deficiency which can be ameliorated by androgen administration. Such compounds are ideal for the treatment of osteoporosis in women and men as a monotherapy or in combination with

5 inhibitors of bone resorption, such as bisphosphonates, estrogens, SERMs, cathepsin K inhibitors,  $\alpha\text{v}\beta 3$  integrin receptor antagonists, calcitonin, and proton pump inhibitors. They can also be used with agents that stimulate bone formation, such as parathyroid hormone or analogs thereof. The SARM compounds of the present invention may also be employed for treatment of prostate disease, such as prostate

10 cancer and benign prostatic hyperplasia (BPH). Moreover, compounds of this invention exhibit minimal effects on skin (acne and facial hair growth) and may be useful for treatment of hirsutism. Additionally, compounds of this invention can stimulate muscle growth and may be useful for treatment of sarcopenia and frailty. They can be employed to reduce subcutaneous and visceral abdominal fat in the

15 treatment of obesity. Moreover, compounds of this invention can exhibit androgen agonism in the central nervous system and may be useful to treat vasomotor symptoms (hot flush) and to increase energy and libido, particularly in postmenopausal women. They may be used as neuroprotective agents in the treatment of Alzheimer's disease. The compounds of the present invention may also be used in

20 the treatment of prostate cancer, either alone or as an adjunct to traditional GnRH agonist/antagonist therapy, for their ability to restore bone, or as a replacement for antiandrogen therapy because of their ability to antagonize androgen in the prostate, and minimize bone depletion in the skeletal system. Further, the compounds of the present invention may be used for their ability to restore bone in the treatment of

25 pancreatic cancer as an adjunct to treatment with antiandrogen, or as monotherapy for their antiandrogenic properties, offering the advantage over traditional antiandrogens of being bone-sparing. Additionally, compounds of this invention can increase the number of blood cells, such as red blood cells and platelets, and may be useful for the treatment of hematopoietic disorders, such as aplastic anemia. Finally, compounds of

30 this invention have minimal effects on lipid metabolism. Thus, considering their

tissue selective androgen receptor agonism listed above, the compounds of this invention are ideal for hormone replacement therapy in hypogonadic (androgen deficient) men.

5 It is therefore an object of the present invention to provide 4-azasteroid derivatives which are useful as selective androgen receptor modulators.

It is another object of the present invention to provide pharmaceutical compositions comprising the 4-azasteroid derivatives of the present invention in association with a pharmaceutically acceptable carrier.

10 It is another object of the present invention to provide pharmaceutical compositions comprising the 4-azasteroid derivatives for use as selective androgen receptor modulators.

It is another object of the present invention to provide methods for the treatment of diseases or conditions caused by androgen deficiency which can be ameliorated by androgen administration.

15 It is another object of the present invention to provide methods for the treatment of diseases or conditions caused by androgen deficiency which can be ameliorated by androgen administration in combination with other agents.


20 It is another object of the present invention to provide 4-azasteroid derivatives and their pharmaceutical compositions for use as a medicament for the treatment of diseases or conditions caused by androgen deficiency which can be ameliorated by androgen administration.

25 It is another object of the present invention to provide 4-azasteroid derivatives and their pharmaceutical compositions for the manufacture of a medicament for the treatment of diseases or conditions caused by androgen deficiency which can be ameliorated by androgen administration.

These and other objects will become readily apparent from the detailed description which follows.

## SUMMARY OF THE INVENTION

The present invention relates to compounds of structural formula I:



or an enantiomer or a pharmaceutically acceptable salt thereof, their uses and

5 phramaceutical composition.

These compounds are effective as androgen receptor agonists and are particularly effective as selective androgen receptor agonists (SARMs). They are therefore useful for the treatment of conditions caused by androgen deficiency or which can be ameliorated by androgen administration.

10           The present invention also relates to pharmaceutical compositions comprising the compounds of the present invention and a pharmaceutically acceptable carrier.


The present invention also relates to methods for the treatment of conditions caused by androgen deficiency or which can be ameliorated by androgen administration in a mammal in need thereof by administering the compounds and pharmaceutical compositions of the present invention.

The present invention also relates to methods for the treatment of osteoporosis, osteopenia, glucocorticoid-induced osteoporosis, periodontal disease, bone fracture, bone damage following bone reconstructive surgery, sarcopenia, frailty, aging skin, male hypogonadism, postmenopausal symptoms in women, atherosclerosis, hypercholesterolemia, hyperlipidemia, obesity, aplastic anemia and other hematopoietic disorders, inflammatory arthritis and joint repair, HIV-wasting, prostate cancer, cancer cachexia, muscular dystrophies, Alzheimer's disease, premature ovarian failure, and autoimmune disease by administering the compounds

and pharmaceutical compositions of the present invention, alone or in combination with a therapeutically effective amount of another agent known to be useful to treat these conditions.

## 5 DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to compounds that are useful as androgen receptor agonists, in particular, as selective androgen receptor agonists. Compounds of the present invention are described by structural formula I:



10

or an enantiomer or a pharmaceutically acceptable salt thereof; wherein

n is 0, 1 or 2;

a-b represents CH=CH or CH<sub>2</sub>CH<sub>2</sub>;

15 "c" represents a double bond;

R<sup>1</sup> is selected from

hydrogen,

hydroxy,

C<sub>1-4</sub> alkoxy,

20 hydroxymethyl, and

C<sub>1-3</sub> alkyl, unsubstituted or substituted with one to seven fluorine atoms;

R<sup>4</sup> is selected from

halogen,

- C<sub>1-6</sub> alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from cyano, carboxy, halogen, hydroxy, oxo, C<sub>1-4</sub> alkoxy, and C<sub>1-4</sub> alkylthio,
- C<sub>2-6</sub> alkenyl,
- 5 C<sub>2-6</sub> alkynyl,
- (CH<sub>2</sub>)<sub>n</sub>-phenyl, and
- (CH<sub>2</sub>)<sub>n</sub>-naphthyl;
- or R<sup>4</sup> together with the carbon atom to which it is attached form a carbonyl or cyclopropyl group, with the proviso that when R<sup>4</sup> together with the
- 10 carbon atom to which it is attached form a carbonyl or cyclopropyl group, then "c" represents a single bond;
- or R<sup>1</sup> and R<sup>4</sup> together with the atoms to which they are attached form a 5- or 6-membered ring system optionally containing an additional heteroatom selected from O, S, and NC<sub>1-4</sub> alkyl;
- 15 R<sup>2</sup> is hydrogen or C<sub>1-4</sub> alkyl;
- R<sup>3</sup> is selected from
- C<sub>1-4</sub> alkyl, wherein said C<sub>1-4</sub> alkyl is optionally substituted with one or more substituents independently selected from halogen, hydroxy, C<sub>1-4</sub> alkoxy, and C<sub>1-4</sub> alkylamino, and
- 20 (CH<sub>2</sub>)<sub>n</sub>-aryl, wherein said aryl is selected from:
- (1) phenyl,
  - (2) naphthyl,
  - (3) benzimidazolyl,
  - (4) benzofuranyl,
  - 25 (5) benzothiophenyl,
  - (6) benzoxazolyl,
  - (7) benzothiazolyl,
  - (8) benzodihydrofuranyl,
  - (9) 1,3-benzodioxolyl,

- (10) 2,3-dihydro-1,4-benzodioxinyl,  
 (11) indolyl,  
 (12) quinolyl,  
 (13) isoquinolyl,  
 5 (14) furanyl,  
 (15) thienyl,  
 (16) imidazolyl,  
 (17) oxazolyl,  
 (18) thiazolyl,  
 10 (19) isoxazolyl,  
 (20) isothiazolyl,  
 (21) pyrazolyl,  
 (22) pyrrolyl,  
 (23) pyridyl,  
 15 (24) pyrimidyl,  
 (25) pyrazinyl,  
 (26) thiadiazolyl,  
 (27) oxadiazolyl,  
 (28) triazolyl,  
 20 (29) imidizopyridinyl,  
 (30) tetrazolyl, and  
 (31) indanyl;

wherein the aryl group as defined in items (1) to (31) is unsubstituted or substituted  
 with one to three groups independently selected from halogen, phenyl,  
 25 C<sub>1-8</sub> alkyl, C<sub>3-8</sub> cycloalkyl, C<sub>3-8</sub> cycloheteroalkyl, phenyl-C<sub>1-6</sub> alkyl,  
 amino-C<sub>0-6</sub> alkyl, C<sub>1-6</sub> alkylamino-C<sub>0-6</sub> alkyl,  
 (C<sub>1-6</sub> alkyl)<sub>2</sub>amino-C<sub>0-6</sub> alkyl, phenyl-C<sub>0-6</sub> alkylamino-C<sub>0-6</sub> alkyl, (phenyl-  
 C<sub>0-6</sub> alkyl)<sub>2</sub>amino-C<sub>0-6</sub> alkyl, C<sub>1-6</sub> alkylthio, phenyl-C<sub>0-6</sub> alkylthio,  
 C<sub>1-6</sub> alkylsulfinyl, phenyl-C<sub>0-6</sub> alkylsulfinyl, C<sub>1-6</sub> alkylsulfonyl,  
 30 phenyl-C<sub>0-6</sub> alkylsulfonyl, C<sub>1-6</sub> alkoxy-C<sub>0-6</sub> alkyl,

phenyl-C<sub>0-6</sub> alkoxy-C<sub>0-6</sub> alkyl, hydroxycarbonyl-C<sub>0-6</sub> alkyl,  
 C<sub>1-6</sub> alkoxycarbonyl-C<sub>0-6</sub> alkyl, phenyl-C<sub>0-6</sub> alkoxycarbonyl-C<sub>0-6</sub> alkyl,  
 hydroxycarbonyl-C<sub>1-6</sub> alkyloxy, hydroxy-C<sub>0-6</sub> alkyl, cyano, nitro,  
 perfluoro-C<sub>1-4</sub> alkyl, perfluoro-C<sub>1-4</sub> alkoxy, C<sub>1-6</sub> alkylcarbonyloxy,  
 5 phenyl-C<sub>0-6</sub> alkylcarbonyloxy, C<sub>1-6</sub> alkylcarbonylamino,  
 phenyl-C<sub>0-6</sub> alkylcarbonylamino, C<sub>1-6</sub> alkylsulfonylamino,  
 phenyl-C<sub>0-6</sub> alkylsulfonylamino, C<sub>1-6</sub> alkoxycarbonylamino,  
 phenyl-C<sub>0-6</sub> alkoxycarbonylamino, C<sub>1-6</sub> alkylaminocarbonylamino,  
 phenyl-C<sub>0-6</sub> alkylaminocarbonylamino, (C<sub>1-6</sub> alkyl)<sub>2</sub> aminocarbonylamino,  
 10 (phenyl-C<sub>0-6</sub> alkyl)<sub>2</sub> aminocarbonylamino, (C<sub>1-6</sub> alkyl)<sub>2</sub> aminocarbonyloxy,  
 and (phenyl-C<sub>0-6</sub> alkyl)<sub>2</sub> aminocarbonyloxy;

the alkyl group is unsubstituted or substituted with one to three substituents  
 independently selected from halogen, hydroxy, and C<sub>1-4</sub> alkoxy;

and wherein any methylene (CH<sub>2</sub>) carbon atom in (CH<sub>2</sub>)<sub>n</sub> is unsubstituted or  
 15 substituted with one to two groups independently selected from halogen,  
 hydroxy, and C<sub>1-4</sub> alkyl; or two substituents when on the same methylene  
 (CH<sub>2</sub>) group are taken together with the carbon atom to which they are  
 attached to form a cyclopropyl group;

or R<sup>2</sup> and R<sup>3</sup> together form a 5- or 6-membered saturated ring fused with a 5- or 6-  
 20 membered aromatic ring system having 0, 1, or 2 heteroatoms selected from  
 N, O, and S.

In one embodiment of the compounds of the present invention are  
 selected from compounds of formula (I) wherein:

25 n is 0, 1 or 2;  
 a-b represents CH=CH or CH<sub>2</sub>CH<sub>2</sub>;  
 "c" represents a double bond;  
 R<sup>1</sup> is selected from  
 hydrogen,

hydroxy,  
C<sub>1-4</sub> alkoxy,  
hydroxymethyl, and  
C<sub>1-3</sub> alkyl, unsubstituted or substituted with one to seven fluorine atoms;

5 R<sup>4</sup> is selected from

halogen,  
C<sub>1-6</sub> alkyl, wherein alkyl is unsubstituted or substituted with one to three  
substituents independently selected from cyano, carboxy, halogen,  
hydroxy, oxo, C<sub>1-4</sub> alkoxy, and C<sub>1-4</sub> alkylthio,

10 C<sub>2-6</sub> alkenyl,  
C<sub>2-6</sub> alkynyl,  
(CH<sub>2</sub>)<sub>n</sub>-phenyl, and  
(CH<sub>2</sub>)<sub>n</sub>-naphthyl;

or R<sup>4</sup> together with the carbon atom to which it is attached form a carbonyl or

15 cyclopropyl group, with the proviso that when R<sup>4</sup> together with the  
carbon atom to which it is attached form a carbonyl or cyclopropyl  
group, then "c" represents a single bond;

or R<sup>1</sup> and R<sup>4</sup> together with the atoms to which they are attached form a 5- or 6-  
membered ring system optionally containing an additional heteroatom selected  
20 from O, S, and NC<sub>1-4</sub> alkyl;

R<sup>2</sup> is hydrogen or C<sub>1-4</sub> alkyl;

R<sup>3</sup> is selected from

C<sub>1-4</sub> alkyl, wherein said C<sub>1-4</sub> alkyl is optionally substituted with one or more  
substituents independently selected from halogen, hydroxy, C<sub>1-4</sub>  
25 alkoxy, and C<sub>1-4</sub> alkylamino, and

(CH<sub>2</sub>)<sub>n</sub>-aryl, wherein said aryl is selected from:

(2) phenyl,  
(2) naphthyl,

- (3) benzimidazolyl,  
(4) benzofuranyl,  
(5) benzothiophenyl,  
(6) benzoxazolyl,  
5 (7) benzothiazolyl,  
(8) benzodihydrofuranyl,  
(9) 1,3-benzodioxolyl,  
(10) 2,3-dihydro-1,4-benzodioxinyl,  
(11) indolyl,  
10 (12) quinolyl,  
(13) isoquinolyl,  
(14) furanyl,  
(15) thienyl,  
(16) imidazolyl,  
15 (17) oxazolyl,  
(18) thiazolyl,  
(19) isoxazolyl,  
(20) isothiazolyl,  
(21) pyrazolyl,  
20 (22) pyrrolyl,  
(23) pyridyl,  
(24) pyrimidyl,  
(25) pyrazinyl,  
(26) thiadiazolyl,  
25 (27) oxadiazolyl,  
(28) triazolyl,  
(29) tetrazolyl, and  
(30) indanyl;

wherein the aryl group as defined in items (1) to (31) is unsubstituted or substituted  
30 with one to three groups independently selected from halogen, phenyl,

C<sub>1-8</sub> alkyl, C<sub>3-8</sub> cycloalkyl, C<sub>3-8</sub> cycloheteroalkyl, phenyl-C<sub>1-6</sub> alkyl,  
 amino-C<sub>0-6</sub> alkyl, C<sub>1-6</sub> alkylamino-C<sub>0-6</sub> alkyl,  
 (C<sub>1-6</sub> alkyl)<sub>2</sub>amino-C<sub>0-6</sub> alkyl, phenyl-C<sub>0-6</sub> alkylamino-C<sub>0-6</sub> alkyl, (phenyl-  
 C<sub>0-6</sub> alkyl)<sub>2</sub>amino-C<sub>0-6</sub> alkyl, C<sub>1-6</sub> alkylthio, phenyl-C<sub>0-6</sub> alkylthio,  
 5 C<sub>1-6</sub> alkylsulfinyl, phenyl-C<sub>0-6</sub> alkylsulfinyl, C<sub>1-6</sub> alkylsulfonyl,  
 phenyl-C<sub>0-6</sub> alkylsulfonyl, C<sub>1-6</sub> alkoxy-C<sub>0-6</sub> alkyl,  
 phenyl-C<sub>0-6</sub> alkoxy-C<sub>0-6</sub> alkyl, hydroxycarbonyl-C<sub>0-6</sub> alkyl,  
 C<sub>1-6</sub> alkoxycarbonyl-C<sub>0-6</sub> alkyl, phenyl-C<sub>0-6</sub> alkoxycarbonyl-C<sub>0-6</sub> alkyl,  
 hydroxycarbonyl-C<sub>1-6</sub> alkyloxy, hydroxy-C<sub>0-6</sub> alkyl, cyano, nitro,  
 10 perfluoro-C<sub>1-4</sub> alkyl, perfluoro-C<sub>1-4</sub> alkoxy, C<sub>1-6</sub> alkylcarbonyloxy,  
 phenyl-C<sub>0-6</sub> alkylcarbonyloxy, C<sub>1-6</sub> alkylcarbonylamino,  
 phenyl-C<sub>0-6</sub> alkylcarbonylamino, C<sub>1-6</sub> alkylsulfonylamino,  
 phenyl-C<sub>0-6</sub> alkylsulfonylamino, C<sub>1-6</sub> alkoxycarbonylamino,  
 phenyl-C<sub>0-6</sub> alkoxycarbonylamino, C<sub>1-6</sub> alkylaminocarbonylamino,  
 15 phenyl-C<sub>0-6</sub> alkylaminocarbonylamino, (C<sub>1-6</sub> alkyl)<sub>2</sub> aminocarbonylamino,  
 (phenyl-C<sub>0-6</sub> alkyl)<sub>2</sub> aminocarbonylamino, (C<sub>1-6</sub> alkyl)<sub>2</sub> aminocarbonyloxy,  
 and (phenyl-C<sub>0-6</sub> alkyl)<sub>2</sub> aminocarbonyloxy;

the alkyl group is unsubstituted or substituted with one to three substituents  
 independently selected from halogen, hydroxy, and C<sub>1-4</sub> alkoxy;

20 and wherein any methylene (CH<sub>2</sub>) carbon atom in (CH<sub>2</sub>)<sub>n</sub> is unsubstituted or  
 substituted with one to two groups independently selected from halogen,  
 hydroxy, and C<sub>1-4</sub> alkyl; or two substituents when on the same methylene  
 (CH<sub>2</sub>) group are taken together with the carbon atom to which they are  
 attached to form a cyclopropyl group;

25 or R<sup>2</sup> and R<sup>3</sup> together form a 5- or 6-membered saturated ring fused with a 5- or 6-  
 membered aromatic ring system having 0, 1, or 2 heteroatoms selected from  
 N, O, and S;

or an enantiomer or a pharmaceutically acceptable salt thereof.

In one embodiment, R<sup>1</sup> is hydrogen or C<sub>1-3</sub> alkyl, unsubstituted or substituted with one to seven fluorine atoms. In another embodiment, R<sup>1</sup> is chosen from hydrogen and methyl.

5                   One variant of the compounds of the present invention are those compounds wherein a-b represents CH<sub>2</sub>CH<sub>2</sub>. In another variant a-b represents CH=CH.

In one embodiment of the invention, R<sup>4</sup> is selected from C<sub>1-4</sub> alkyl, such as for example, methyl.

10                   In another embodiment, R<sup>4</sup> together with the carbon atom to which it is attached form a carbonyl or cyclopropyl group. In one variant, the group is a cyclopropyl group.

In yet another embodiment of the invention, R<sup>1</sup> and R<sup>4</sup> together with the atoms to which they are attached form a 5- or 6-membered ring system optionally  
15                   containing an additional heteroatom selected from O, S, and NC<sub>1-4</sub> alkyl.

In one embodiment of the invention, R<sup>2</sup> is hydrogen.

In yet another embodiment, R<sup>3</sup> is (CH<sub>2</sub>)<sub>n</sub>-aryl. In one class of this embodiment, (CH<sub>2</sub>)<sub>n</sub>-aryl and n is selected from 0 and 1, and aryl is selected from:  
20                   phenyl, imidazopyridinyl, benzimidazolyl, benzothiophenyl, indolyl, quinolyl, isoquinolyl, thienyl, imidazolyl, thiazolyl, pyrazolyl, pyrrolyl, pyridyl, pyrimidyl, pyrazinyl, thiadiazolyl, triazolyl, and indanyl. In a class of this embodiment, R<sup>3</sup> is selected from phenyl, benzimidazolyl, benzothiophenyl, indolyl, pyrazolyl, pyrrolyl, pyridyl, pyrimidyl, pyrazinyl, thiadiazolyl, imidazopyridinyl, and triazolyl.

In another embodiment, R<sup>1</sup> is methyl and R<sup>3</sup> is selected from phenyl,  
25                   benzimidazolyl, benzothiophenyl, indolyl, pyrazolyl, pyrrolyl, pyridyl, pyrimidyl, pyrazinyl, thiadiazolyl, imidazopyridinyl, and triazolyl. In class of this embodiment, R<sup>4</sup> is selected from C<sub>1-4</sub> alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from cyano, carboxy, halogen, hydroxy, oxo, C<sub>1-4</sub> alkoxy, and C<sub>1-4</sub> alkylthio, (CH<sub>2</sub>)<sub>n</sub>-phenyl, and halogen.

In one embodiment of the invention, R<sup>4</sup> is a methyl group. In another embodiment, R<sup>4</sup> together with the carbon atom to which it is attached form a cyclopropyl group.

Illustrative but nonlimiting examples of compounds of the present invention that are useful as androgen receptor modulators are the following:

5 N-phenyl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(3-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
10 carboxamide,  
N-(4-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(2-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
15 N-(3-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(2-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
20 carboxamide,  
N-(3-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(4-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
25 N-(2,2,2-trifluoroethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(2-fluorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(3-fluorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
30 carboxamide,

- N-(4-fluorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-(2,3-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,5-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-(2-chloro-6-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-chloro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(4-chloro-2-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-(2,4-dichloro-5-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[4-(morpholin-4-yl)phenyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 20 N-(4-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-[2-(piperidin-1-yl)phenyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(5-fluoro-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 30

- N-(4-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(5-methyl-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- 5 N-(4-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(3-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
10 carboxamide,
- N-(3-cyanophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-(pyridin-3-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2-chloro-4-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2,6-dichlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
20 carboxamide,
- N-(2,6-difluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2,3,4,6-tetrafluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- 25 N-(2,3-dichlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2,5-dichlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

- N-(4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(3-chloro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- 5 N-(3-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(3-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(4-acetyl-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
10 carboxamide,
- N-(4-methyl-thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(4-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- 15 N-(2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2,5-difluoro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-  
17 $\beta$ -carboxamide,
- N-(2,4,5,6-tetrafluoro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-  
20 en-17 $\beta$ -carboxamide,
- N-(2,6-dichloro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-  
17 $\beta$ -carboxamide,
- N-(3,4-dicyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- 25 N-(2,4-dichloro-6-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-  
17 $\beta$ -carboxamide,
- N-(2-chloro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(3-fluoro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
30 carboxamide,

- N-(3-trifluoromethyl-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-chloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-(3,5-dichloro-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-chloro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,5-dichloro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-(3-chloro-5-trifluoromethyl-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,6-dichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 3-{[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino}-4-chlorobenzoic acid,
- N-(4-chlorobenzothiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(benzothiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 20 N-(4,5-dimethylthiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,3,5-trichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-(2,6-dichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(5-ethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(4-methyl-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 30

- N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-ehloro-6-trifluoromethyl-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-(2,6-dichloro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,6-dichloro-6'-trifluoro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-(4-methyl-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(pyrazin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(6-chloro-pyrazin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-(benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-fluoro-6-chlorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 20 N-((R)-phenylmethylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-((S)-phenylmethylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,6-dichlorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-(3,4-dichlorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[(1S,2S)-2-hydroxycyclohexyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

- N-(3H-imidazo[4,5-b]pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[(1R,2S)-2-hydroxycyclohexyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-(1,3-thiazol-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(1H-indol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[2-(1H-benzimidazol-2-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-[2-(6-chloro-1H-benzimidazol-2-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- tert-butyl 3-({[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino}methyl)-1H-indole-1-carboxylate,
- 15 tert-butyl 3-(2-{[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino}ethyl)-1H-indole-1-carboxylate,
- N-(benzimidazol-2-ylmethylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 20 N-[2-(1-methyl-1H-benzimidazol-2-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(imidazo[1,2-a]pyrimidin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-(1H-indol-5-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[(6-chloro-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(imidazo[1,2-a]pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 30

- N-(2-imidazo[1,2-a]pyridin-2-ylethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-[2-(2-phenyl-1H-indol-3-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
5 N-(1H-indol-7-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(1H-indol-4-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(imidazo[1,2-a]pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
10 N-[(1-phenyl-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(pyrazolo[1,5-a]pyridin-7-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
15 N-(pyrazolo[1,5-a]pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-[(1-methyl-2,3-dihydro-1H-indol-5-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-piperidin-4-yl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
20 N-piperidin-3-yl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2-ethylpyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-[(1-methyl-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
25 N-(5-t-Butyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(5-trifluoromethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2-methoxypyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
30

- N-(2-ethoxypyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(4-trifluoromethylpyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-  
5 17 $\beta$ -carboxamide,  
N-(4,6-dimethylpyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(6-methylpyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
10 N-(7H-purin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(5-cyclopropyl-1H-pyrazol-3-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(6-methoxybenzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-  
15 5-en-17 $\beta$ -carboxamide,  
N-(2,3-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2,6-dichlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
20 N-(2-fluoro-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-fluoro-6-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-trifluoromethyl-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-fluoro-4-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
25 N-(2-trifluoromethylphenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-fluoro-5-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-phenyl-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(5-ethyl-1,3,4-thiadiazol-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

- N-(5-methyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 5 N-(pyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(2-chloropyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(2-methylpyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(2-chloropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(4-fluoropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 10 N-(4-chloropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(4-methylpyridin-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(4-chloropyridin-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(benzimidazol-2-ylmethyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 15 N-((N-methylbenzimidazol-2-yl)methyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(3H-imidazo[4,5-b]pyridin-2-ylmethyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(pyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 20 N-(2,3-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(2-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(2,6-dichlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 25 N-(2-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(2-fluoro-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(2,6-difluorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,

- N-(2-trifluoromethyl-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(2-fluoro-4-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 5 N-(2-trifluoromethylphenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(2-fluoro-5-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-phenyl-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 10 N-(2,6-dichloro-4-cyanophenyl) -6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(1-methyl-6-trifluoromethyl-1H-benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(2-methyl-3-chloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 15 enantiomers, and pharmaceutically acceptable salts thereof.

In one embodiment, the compounds of the invention are selected from:

- 20 N-(3H-imidazo[4,5-b]pyridin-2-ylmethyl) -4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(imidazo[1,2-a]pyrimidin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(imidazo[1,2-a]pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 25 N-(2-imidazo[1,2-a]pyridin-2-ylethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(3H-imidazo[4,5-b]pyridin-2-ylmethyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 30 enantiomers, and pharmaceutically acceptable salts thereof.

In a class of this embodiment, the compounds are selected from:

N-(3H-imidazo[4,5-b]pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

5 N-(imidazo[1,2-a]pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

N-(3H-imidazo[4,5-b]pyridin-2-ylmethyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

enantiomers, and pharmaceutically acceptable salts thereof.

10 In yet another embodiment of the invention, the compounds of formula

(I) are selected from:

N-phenyl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

N-(2-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

15 N-(3-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

N-(4-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

20 N-(2-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

N-(3-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

N-(4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

25 N-(2-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

N-(3-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

30 N-(4-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

- N-(2,2,2-trifluoroethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2-fluorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- 5 N-(3-fluorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(4-fluorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
10 carboxamide,
- N-(2,3-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-  
5-en-17 $\beta$ -carboxamide,
- N-(2,5-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-  
5-en-17 $\beta$ -carboxamide,
- 15 N-(2-chloro-6-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-  
en-17 $\beta$ -carboxamide,
- N-(2-chloro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-  
en-17 $\beta$ -carboxamide,
- N-(4-chloro-2-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-  
20 en-17 $\beta$ -carboxamide,
- N-(2,4-dichloro-5-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-  
17 $\beta$ -carboxamide,
- N-[4-(morpholin-4-yl)phenyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- 25 N-(3-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(4-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
30 carboxamide,

- N-[2-(piperidin-1-yl)phenyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(5-fluoro-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-(2-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(4-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(5-methyl-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-(4-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-(3-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3-cyanophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 20 N-(pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(pyridin-3-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-chloro-4-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-(2,6-dichlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,6-difluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,3,4,6-tetrafluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 30

- N-(2,3-dichlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,5-dichlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-(thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3-chloro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-(3-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(3-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(4-acetyl-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-(4-methyl-thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(4-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 20 N-(2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2,5-difluoro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,4,5,6-tetrafluoro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-(2,6-dichloro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(3,4-dicyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

- N-(2,4-dichloro-6-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-chloro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-(3-fluoro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3-trifluoromethyl-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-chloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-(3,5-dichloro-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-chloro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-(2,5-dichloro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3-chloro-5-trifluoromethyl-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,6-dichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 20 3-{[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino}-4-chlorobenzoic acid,
- N-(4-chlorobenzothiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-(benzothiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(4,5-dimethylthiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,3,5-trichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 30

- N-(2,6-dichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(5-ethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-(4-methyl-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-chloro-6-trifluoromethyl-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-(2,6-dichloro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,6-dichloro-6'-trifluoro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(4-methyl-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(pyrazin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 20 N-(6-chloro-pyrazin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-fluoro-6-chlorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-((R)-phenylmethylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-((S)-phenylmethylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

- N-(2,6-dichlorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3,4-dichlorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-[(1S,2S)-2-hydroxycyclohexyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[(1R,2S)-2-hydroxycyclohexyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(1,3-thiazol-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-(1H-indol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[2-(1H-benzimidazol-2-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-[2-(6-chloro-1H-benzimidazol-2-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- tert-butyl 3-({[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino}methyl)-1H-indole-1-carboxylate,
- 20 tert-butyl 3-(2-{[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino}ethyl)-1H-indole-1-carboxylate,
- N-(benzimidazol-2-ylmethylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-[2-(1-methyl-1H-benzimidazol-2-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(1H-indol-5-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[(6-chloro-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 30

- N-[2-(2-phenyl-1H-indol-3-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(1H-indol-7-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-(1H-indol-4-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[(1-phenyl-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(pyrazolo[1,5-a]pyridin-7-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-(pyrazolo[1,5-a]pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[(1-methyl-2,3-dihydro-1H-indol-5-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-piperidin-4-yl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-piperidin-3-yl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-ethylpyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[(1-methyl-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 20 N-(5-t-Butyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(5-trifluoromethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-(2-methoxypyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-ethoxypyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

- N-(4-trifluoromethylpyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(4,6-dimethylpyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-(6-methylpyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(7H-purin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(5-cyclopropyl-1H-pyrazol-3-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-(6-methoxybenzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(1-methyl-6-trifluoromethyl-1H-benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-(2,3-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2,6-dichlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2-fluoro-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 20 N-(2-fluoro-6-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2-trifluoromethyl-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2-fluoro-4-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2-trifluoromethylphenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 25 N-(2-fluoro-5-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-phenyl-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(5-ethyl-1,3,4-thiadiazol-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(5-methyl-1,3,4-thiadiazol-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 30

- N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(pyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2-chloropyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 5 N-(2-methylpyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2-chloropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(4-fluoropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(4-chloropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(4-methylpyridin-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 10 N-(4-chloropyridin-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(benzimidazol-2-ylmethyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-((N-methylbenzimidazol-2-yl)methyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 15 N-(pyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2,3-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2,6-dichlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 20 N-(2-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2-fluoro-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2,6-difluorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 25 N-(2-trifluoromethyl-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2-fluoro-4-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,

- N-(2-trifluoromethylphenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-fluoro-5-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
5 N-phenyl-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2,6-dichloro-4-cyanophenyl) -6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-methyl-3-chloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
10 enantiomers, and pharmaceutically acceptable salts thereof.

In one variant, the compounds of formula (I) are selected from:

- N-phenyl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(3-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
15 carboxamide,  
N-(3-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(3-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
20 N-(2,3-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2,5-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(3-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
25 carboxamide,  
N-(2-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(3-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

- N-(3-cyanophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2,6-difluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- 5 N-(2,3,4,6-tetrafluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(3-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- 10 N-(4-methyl-thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(4-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
15 carboxamide,
- N-(2-chloro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(3-fluoro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- 20 N-(2-chloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(3,5-dichloro-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(4,5-dimethylthiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
25 carboxamide,
- N-(2,3,5-trichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2,6-dichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,

- N-(5-ethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(4-methyl-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(6-chloro-pyrazin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-(benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[(1S,2S)-2-hydroxycyclohexyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-[(1R,2S)-2-hydroxycyclohexyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- tert-butyl 3-({[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino)methyl)-1H-indole-1-carboxylate,
- 20 N-[(6-chloro-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(1-methyl-6-trifluoromethyl-1H-benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-ethylpyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-[(1-methyl-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(5-t-Butyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

- N-(5-trifluoromethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2-methoxypyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
5 N-(2-methyl-3-chloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
enantiomers, and pharmaceutically acceptable salts thereof.

- In one class of this variant, the compounds of the present invention are  
10 selected from:  
N-phenyl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(3-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(3-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
15 carboxamide,  
N-(3-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2,3-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
20 N-(2,5-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(3-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
25 carboxamide,  
N-(3-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(3-cyanophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

- N-(2,6-difluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,3,4,6-tetrafluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-(3-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(4-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -
- 10 carboxamide,
- N-(2-chloro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3-fluoro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-(benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[(1S,2S)-2-hydroxycyclohexyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[(1R,2S)-2-hydroxycyclohexyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-
- 20 17 $\beta$ -carboxamide,
- tert-butyl 3-({[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino}methyl)-1H-indole-1-carboxylate,
- N-[(6-chloro-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -
- 25 androst-5-en-17 $\beta$ -carboxamide,
- N-(1-methyl-6-trifluoromethyl-1H-benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[(1-methyl-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -
- androst-5-en-17 $\beta$ -carboxamide,
- 30 enantiomers, and pharmaceutically acceptable salts thereof.

In yet another class, the compounds of the invention are selected from:

N-(thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

N-(4-methyl-thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -

5 carboxamide,

N-(4,5-dimethylthiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -

carboxamide,

N-(5-ethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-

17 $\beta$ -carboxamide,

10 N-(5-t-Butyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-

17 $\beta$ -carboxamide,

N-(5-trifluoromethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -

androst-5-en-17 $\beta$ -carboxamide,

N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-

15 5-en-17 $\beta$ -carboxamide,

N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-

17 $\beta$ -carboxamide,

enantiomers, and pharmaceutically acceptable salts thereof.

20 In yet another embodiment, the compounds of the invention are selected from:

N-(2-chloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -

carboxamide,

N-(3,5-dichloro-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -

25 carboxamide,

N-(2,3,5-trichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -

carboxamide,

N-(2,6-dichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -

carboxamide,

N-(4-methyl-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,

N-(6-chloro-pyrazin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,

5 N-(2-ethylpyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,

N-(2-methoxypyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,

enantiomers, and pharmaceutically acceptable salts thereof.

10

In still another embodiment, the compounds of formula (I) are selected  
from:

N-(2,3-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-(2-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

15 N-(2,6-dichlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-(2-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-(2-fluoro-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-(2-fluoro-6-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-(2-trifluoromethyl-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -

20 carboxamide,

N-(2-fluoro-4-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-(2-trifluoromethylphenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-(2-fluoro-5-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-phenyl-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

25 N-(5-ethyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,

N-(5-methyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,

N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -

30 carboxamide,

- N-(pyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-chloropyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-methylpyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-chloropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
5 N-(4-fluoropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(4-chloropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(4-methylpyridin-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(4-chloropyridin-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(benzimidazol-2-ylmethyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
10 carboxamide,  
N-((N-methylbenzimidazol-2-yl)methyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
N-(pyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2,3-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -  
15 carboxamide,  
N-(2-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2,6-dichlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
N-(2-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
20 N-(2-fluoro-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
N-(2,6-difluorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
N-(2-trifluoromethyl-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -  
25 androst-17 $\beta$ -carboxamide,  
N-(2-fluoro-4-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
N-(2-trifluoromethylphenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,

N-(2-fluoro-5-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-phenyl-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2,6-dichloro-4-cyanophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-  
5 17 $\beta$ -carboxamide,  
enantiomers, and pharmaceutically acceptable salts thereof.

In a class of this embodiment, the compounds are selected from:

N-(5-ethyl-1,3,4-thiadiazol-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
10 carboxamide,  
N-(5-methyl-1,3,4-thiadiazol-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
15 enantiomers, and pharmaceutically acceptable salts thereof.

In another class of this embodiment, the compounds of formula (1) are selected from:

N-(pyridin-4-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
20 N-(2-chloropyridin-4-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-methylpyridin-4-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-chloropyridin-3-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(4-fluoropyridin-3-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(4-chloropyridin-3-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
25 N-(4-methylpyridin-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(4-chloropyridin-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(pyridin-3-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
enantiomers, and pharmaceutically acceptable salts thereof.

In yet another class of this embodiment, the compounds of the present invention are selected from:

N-(2,3-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-(4-methylpyridin-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

5 N-(2,3-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-(2-fluoro-5-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

enantiomers, and pharmaceutically acceptable salts thereof.

10

The term "alkyl" shall mean straight or branched chain alkanes of one to ten total carbon atoms, or any number within this range (i.e., methyl, ethyl, 1-propyl, 2-propyl, n-butyl, s-butyl, t-butyl, etc.). The term "C<sub>0</sub> alkyl" (as in "C<sub>0</sub>-8 alkylaryl") shall refer to the absence of an alkyl group.

15

The term "alkenyl" shall mean straight or branched chain alkenes of two to ten total carbon atoms, or any number within this range.

The term "alkynyl" shall mean straight or branched chain alkynes of two to ten total carbon atoms, or any number within this range.

20 The term "alkylidene" shall mean a straight or branched chain alkylidene group of one to ten total carbon atoms, or any number within this range.

The term "cycloalkyl" shall mean cyclic rings of alkanes of three to eight total carbon atoms, or any number within this range (i.e., cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, or cyclooctyl).

25 The term "cycloheteroalkyl," as used herein, shall mean a 3- to 8-membered fully saturated heterocyclic ring containing one or two heteroatoms chosen from N, O, or S. Examples of cycloheteroalkyl groups include, but are not limited to, piperidinyl, pyrrolidinyl, azetidiny, morpholinyl, and piperazinyl. In one embodiment of the present invention, cycloheteroalkyl is selected from piperidinyl, pyrrolidinyl, and morpholinyl.

The term "alkoxy," as used herein, refers to straight or branched chain alkoxides of the number of carbon atoms specified (e.g., C<sub>1-5</sub> alkoxy), or any number within this range (i.e., methoxy, ethoxy, etc.).

The term "aryl," as used herein, refers to a monocyclic or bicyclic system comprising at least one aromatic ring, wherein the monocyclic or bicyclic system contains 0, 1, 2, 3, or 4 heteroatoms chosen from N, O, or S, and wherein the monocyclic or bicyclic system is either unsubstituted or substituted with one or more groups independently selected from halogen, phenyl, C<sub>1-8</sub> alkyl, C<sub>3-8</sub> cycloalkyl, C<sub>3-8</sub> cycloheteroalkyl, phenyl-C<sub>1-6</sub> alkyl, amino-C<sub>0-6</sub> alkyl, C<sub>1-6</sub> alkylamino-C<sub>0-6</sub> alkyl, (C<sub>1-6</sub> alkyl)<sub>2</sub>amino-C<sub>0-6</sub>alkyl, phenyl-C<sub>0-6</sub> alkylamino-C<sub>0-6</sub>alkyl, (phenyl-C<sub>0-6</sub> alkyl)<sub>2</sub>amino-C<sub>0-6</sub> alkyl, C<sub>1-6</sub> alkylthio, phenyl-C<sub>0-6</sub> alkylthio, C<sub>1-6</sub> alkylsulfinyl, phenyl-C<sub>0-6</sub> alkylsulfinyl, C<sub>1-6</sub> alkylsulfonyl, phenyl-C<sub>0-6</sub> alkylsulfonyl, C<sub>1-6</sub> alkoxy-C<sub>0-6</sub>alkyl, phenyl-C<sub>0-6</sub> alkoxy-C<sub>0-6</sub> alkyl, hydroxycarbonyl-C<sub>0-6</sub> alkyl, C<sub>1-6</sub> alkoxycarbonyl-C<sub>0-6</sub> alkyl, phenyl-C<sub>0-6</sub> alkoxycarbonyl-C<sub>0-6</sub>alkyl, hydroxycarbonyl C<sub>1-6</sub> alkyloxy, hydroxy-C<sub>0-6</sub> alkyl, cyano, nitro, perfluoroC<sub>1-4</sub> alkyl, perfluoroC<sub>1-4</sub> alkoxy, oxo, C<sub>1-6</sub> alkylcarbonyloxy, phenyl-C<sub>0-6</sub> alkylcarbonyloxy, C<sub>1-6</sub> alkylcarbonylamino, phenyl-C<sub>0-6</sub> alkylcarbonylamino, C<sub>1-6</sub> alkylsulfonylamino, phenyl-C<sub>0-6</sub> alkylsulfonylamino, C<sub>1-6</sub> alkoxycarbonylamino, phenyl-C<sub>0-6</sub> alkoxycarbonylamino, C<sub>1-6</sub> alkylaminocarbonylamino, phenyl-C<sub>0-6</sub> alkylaminocarbonylamino, (C<sub>1-6</sub> alkyl)<sub>2</sub> aminocarbonylamino, (phenyl-C<sub>0-6</sub> alkyl)<sub>2</sub> aminocarbonylamino, (C<sub>1-6</sub> alkyl)<sub>2</sub> aminocarbonyloxy, and (phenyl-C<sub>0-6</sub> alkyl)<sub>2</sub> aminocarbonyloxy. Preferably, the aryl group is unsubstituted, mono-, di-, or tri-substituted with one to three of the above-named substituents; more preferably, the aryl group is unsubstituted, mono- or di-substituted with one to two of the above-named substituents.

Whenever the term "alkyl" or "aryl" or either of their prefix roots appears in a name of a substituent (e.g., aryl C<sub>0-8</sub> alkyl), it shall be interpreted as including those limitations given above for "alkyl" and "aryl." Designated numbers of carbon atoms (e.g., C<sub>0-8</sub>) shall refer independently to the number of carbon atoms in

an alkyl or cyclic alkyl moiety or to the alkyl portion of a larger substituent in which alkyl appears as its prefix root.

The terms "arylalkyl" and "alkylaryl" include an alkyl portion where alkyl is as defined above and include an aryl portion where aryl is as defined above.

5 Examples of arylalkyl include, but are not limited to, benzyl, fluorobenzyl, chlorobenzyl, phenylethyl, phenylpropyl, fluorophenylethyl, chlorophenylethyl, thienylmethyl, thienylethyl, and thienylpropyl. Examples of alkylaryl include, but are not limited to, toluene, ethylbenzene, propylbenzene, methylpyridine, ethylpyridine, propylpyridine and butylpyridine.


10 The term "halogen" shall include iodine, bromine, chlorine, and fluorine.

The term "oxy" means an oxygen (O) atom. The term "thio" means a sulfur (S) atom. The term "oxo" means " $=O$ ". The term "carbonyl" means " $C=O$ ".

The term "substituted" shall be deemed to include multiple degrees of  
15 substitution by a named substituent. Where multiple substituent moieties are disclosed or claimed, the substituted compound can be independently substituted by one or more of the disclosed or claimed substituent moieties, singly or plurally. By independently substituted, it is meant that the (two or more) substituents can be the same or different.

20 When any variable (e.g.,  $R^2$ ,  $R^3$ , etc.) occurs more than one time in any substituent or in formula I, its definition in each occurrence is independent of its definition at every other occurrence. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

Under standard nomenclature used throughout this disclosure, the  
25 terminal portion of the designated side chain is described first, followed by the adjacent functionality toward the point of attachment. For example, a  $C_{1-5}$  alkylcarbonylamino  $C_{1-6}$  alkyl substituent is equivalent to



In choosing compounds of the present invention, one of ordinary skill in the art will recognize that the various substituents, i.e. R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, etc., are to be chosen in conformity with well-known principles of chemical structure connectivity.

Compounds of the present invention have been found to be tissue-selective modulators of the androgen receptor (SARMs). In one aspect, compounds of the present invention may be useful to activate the function of the androgen receptor in a mammal, and in particular to activate the function of the androgen receptor in bone and/or muscle tissue and block or inhibit ("antagonize") the function of the androgen receptor in the prostate of a male individual or in the uterus of a female individual. The activation of the AR in bone can be assayed through stimulation of bone formation in a rodent model of osteoporosis, and the antagonism of the AR in the prostate can be assayed through observation of minimal effects on prostate growth in castrated rodents and antagonism of prostate growth induced by AR agonists, as detailed in the Examples.

A further aspect of the present invention is concerned with compounds of structural formula I that block the function of the androgen receptor in the prostate of a male individual or in the uterus of a female individual induced by AR agonists, but not in hair-growing skin or vocal cords, and activate the function of the androgen receptor in bone and/or muscle tissue, but not in organs which control blood lipid levels (e.g. liver). The compounds of the present invention may be used to treat conditions which are caused by androgen deficiency or which can be ameliorated by androgen replacement, including, but not limited to osteoporosis, osteopenia, glucocorticoid-induced osteoporosis, periodontal disease, bone fracture, bone damage following bone reconstructive surgery, sarcopenia, frailty, aging skin, male hypogonadism, postmenopausal symptoms in women, atherosclerosis, hypercholesterolemia, hyperlipidemia, obesity, aplastic anemia and other hematopoietic disorders, inflammatory arthritis and joint repair, HIV-wasting, Alzheimer's disease, prostate cancer, cancer cachexia, muscular dystrophies, Alzheimer's disease, premature ovarian failure, and autoimmune disease, alone or in combination with other active agents. Treatment is effected by administration of a

therapeutically effective amount of a compound of structural formula I to a mammal in need of such treatment. In addition, these compounds are useful as ingredients in pharmaceutical compositions alone or in combination with other active agents.

In one embodiment, the compounds of the present invention may be used to treat conditions in a male individual which are caused by androgen deficiency or which can be ameliorated by androgen replacement, including, but not limited to, osteoporosis, osteopenia, glucocorticoid-induced osteoporosis, periodontal disease, HIV-wasting, prostate cancer, cancer cachexia, obesity, aplastic and other anemias, muscular dystrophies, and Alzheimer's disease, alone or in combination with other active agents. Treatment is effected by administration of a therapeutically effective amount of a compound of structural formula I to a male individual in need of such treatment.

In another embodiment, the compounds of the present invention may be used to treat conditions in a female individual which are caused by androgen deficiency or which can be ameliorated by androgen replacement, including, but not limited to, osteoporosis, osteopenia, glucocorticoid-induced osteoporosis, postmenopausal symptoms, periodontal disease, HIV-wasting, cancer cachexia, obesity, aplastic and other anemias, muscular dystrophies, Alzheimer's disease, premature ovarian failure, and autoimmune disease, alone or in combination with other active agents. Treatment is effected by administration of a therapeutically effective amount of a compound of structural formula I to a female individual in need of such treatment.

The compounds of structural formula I may also be employed as adjuncts to traditional androgen depletion therapy in the treatment of prostate cancer to restore bone, minimize bone loss, and maintain bone mineral density. In this manner, they may be employed together with traditional androgen deprivation therapy, including GnRH agonists/antagonists, such as those disclosed in P. Limonta, et al., "LHRH analogues as anticancer agents: pituitary and extrapituitary sites of action," Exp. Opin. Invest. Drugs, 10: 709-720 (2001); H.J. Stricker, "Luteinizing hormone-releasing hormone antagonists," Urology, 58 (Suppl. 2A): 24-27 (2001);

R.P. Millar, et al., "Progress towards the development of non-peptide orally-active GnRH antagonists," British Medical Bulletin, 56: 761-772 (2000); and A.V. Schally et al., "Rational use of agonists and antagonists of LH-RH in the treatment of hormone-sensitive neoplasms and gynecologic conditions," Advanced Drug Delivery Reviews, 28: 157-169 (1997). It is also possible that the compounds of structural formula I may be used in combination with antiandrogens, such as flutamide, 2-hydroxyflutamide (the active metabolite of flutamide), nilutamide, and bicalutamide (Casodex™) in the treatment of prostate cancer.

Further, the compounds of the present invention may also be employed in the treatment of pancreatic cancer, either for their androgen antagonist properties or as an adjunct to an antiandrogen, such as flutamide, 2-hydroxyflutamide (the active metabolite of flutamide), nilutamide, and bicalutamide (Casodex™).

Compounds of structural formula I have minimal negative effects on lipid metabolism. Therefore, considering their tissue selective androgen agonistic properties, the compounds of this invention have advantages over existing approaches for hormone replacement therapy in hypogonadic (androgen deficient) male individuals.

Additionally, compounds of the present invention can increase the number of blood cells, such as red blood cells and platelets, and can be used for treatment of hematopoietic disorders, such as aplastic anemia.

Representative compounds of the present invention typically display submicromolar binding affinity for the androgen receptor. Compounds of this invention are therefore useful in treating mammals suffering from disorders related to androgen receptor function. Pharmacologically effective amounts of the compound, including the pharmaceutically effective salts thereof, are administered to the mammal, to treat disorders related to androgen receptor function, or which can be improved by the addition of additional androgen, such as osteoporosis, periodontal disease, bone fracture, bone damage following bone reconstructive surgery, sarcopenia, frailty, aging skin, male hypogonadism, postmenopausal symptoms in women, atherosclerosis, hypercholesterolemia, hyperlipidemia, obesity, aplastic

anemia and other hematopoietic disorders, pancreatic cancer, Alzheimer's disease, inflammatory arthritis, and joint repair.

It is generally preferable to administer compounds of the present invention in their enantiomerically pure form. Racemic mixtures can be separated  
5 into their individual enantiomers by any of a number of conventional methods. These include chiral chromatography, derivatization with a chiral auxiliary followed by separation by chromatography or crystallization, and fractional crystallization of diastereomeric salts.

As used herein, a compound of the present invention which functions  
10 as an "agonist" of the androgen receptor can bind to the androgen receptor and initiate a physiological or a pharmacological response characteristic of that receptor. The term "tissue-selective androgen receptor modulator" refers to an androgen receptor ligand that mimics the action of a natural ligand in some tissues but not in others. A "partial agonist" is an agonist which is unable to induce maximal activation of the  
15 receptor population, regardless of the amount of compound applied. A "full agonist" induces full activation of the androgen receptor population at a given concentration. A compound of the present invention which functions as an "antagonist" of the androgen receptor can bind to the androgen receptor and block or inhibit the androgen-associated responses normally induced by a natural androgen receptor  
20 ligand.

The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic or organic bases and inorganic or organic acids. Salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium,  
25 manganic salts, manganous, potassium, sodium, zinc, and the like. Particularly preferred are the ammonium, calcium, lithium, magnesium, potassium, and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, and basic ion exchange resins,  
30 such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine,

diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethyl-morpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine, and the like.

When the compound of the present invention is basic, salts may be prepared from pharmaceutically acceptable non-toxic acids, including inorganic and organic acids. Such acids include acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, formic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, malonic, mucic, nitric, pantoic, pantothenic, phosphoric, propionic, succinic, sulfuric, tartaric, p-toluenesulfonic acid, trifluoroacetic acid, and the like. Particularly preferred are citric, fumaric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, and tartaric acids.

The term "therapeutically effective amount" means the amount the compound of structural formula I that will elicit the biological or medical response of a tissue, system, animal or human that is being sought by the researcher, veterinarian, medical doctor or other clinician.

The term "composition" as used herein is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

By "pharmaceutically acceptable" it is meant that the carrier, diluent or excipient must be compatible with the other ingredients of the formulation and not be deleterious to the recipient thereof.

The terms "administration of a compound" and "administering a compound" should be understood to mean providing a compound of the invention or a prodrug of a compound of the invention to the individual in need of treatment.

By the term "modulating a function mediated by the androgen receptor in a tissue selective manner" is meant modulating a function mediated by the

androgen receptor selectively (or discriminately) in anabolic (bone and/or muscular) tissue (bone and muscular) in the absence of such modulation at androgenic (reproductive) tissue, such as the prostate, testis, seminal vesicles, ovary, uterus, and other sex accessory tissues. In one embodiment the function of the androgen receptor  
5 in anabolic tissue is activated whereas the function of the androgen receptor in androgenic tissue is blocked or suppressed.

The administration of a compound of structural formula I in order to practice the present methods of therapy is carried out by administering an effective amount of the compound of structural formula I to the patient in need of such  
10 treatment or prophylaxis. The need for a prophylactic administration according to the methods of the present invention is determined via the use of well-known risk factors. The effective amount of an individual compound is determined, in the final analysis, by the physician in charge of the case, but depends on factors such as the exact disease to be treated, the severity of the disease and other diseases or conditions from which  
15 the patient suffers, the chosen route of administration, other drugs and treatments which the patient may concomitantly require, and other factors in the physician's judgment.

Generally, the daily dosage of a compound of structural formula I may be varied over a wide range from 0.01 to 1000 mg per adult human per day. Most  
20 preferably, dosages range from 0.1 to 200 mg/day. For oral administration, the compositions are preferably provided in the form of tablets containing 0.01 to 1000 mg, particularly 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 3.0, 5.0, 6.0, 10.0, 15.0, 25.0, 50.0, 75, 100, 125, 150, 175, 180, 200, 225, and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the mammal to be treated.

25 The dose may be administered in a single daily dose or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, based on the properties of the individual compound selected for administration, the dose may be administered less frequently, e.g., weekly, twice weekly, monthly, etc. The unit dosage will, of course, be correspondingly larger for  
30 the less frequent administration.

When administered via intranasal routes, transdermal routes, by rectal or vaginal suppositories, or through an intravenous solution, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

Exemplifying the invention is a pharmaceutical composition comprising any of the compounds described above and a pharmaceutically acceptable carrier. Also exemplifying the invention is a pharmaceutical composition made by combining any of the compounds described above and a pharmaceutically acceptable carrier. An illustration of the invention is a process for making a pharmaceutical composition comprising combining any of the compounds described above and a pharmaceutically acceptable carrier.

Formulations of the tissue-selective androgen receptor modulator employed in the present method for medical use comprise a compound of structural formula I together with an acceptable carrier thereof and optionally other therapeutically active ingredients. The carrier must be pharmaceutically acceptable in the sense of being compatible with the other ingredients of the formulation and not being deleterious to the recipient subject of the formulation.

The present invention, therefore, further provides a pharmaceutical formulation comprising a compound of structural formula I together with a pharmaceutically acceptable carrier thereof.

The formulations include those suitable for oral, rectal, intravaginal, topical or parenteral (including subcutaneous, intramuscular and intravenous administration). Preferred formulations are those suitable for oral administration.

The formulations may be presented in a unit dosage form and may be prepared by any of the methods known in the art of pharmacy. All methods include the step of bringing the active compound in association with a carrier which constitutes one or more ingredients. In general, the formulations are prepared by uniformly and intimately bringing the active compound in association with a liquid carrier, a waxy solid carrier or a finely divided solid carrier, and then, if needed, shaping the product into the desired dosage form.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets, tablets or lozenges, each containing a predetermined amount of the active compound; as a powder or granules; or a suspension or solution in an aqueous liquid or non-aqueous liquid, e.g., a syrup,  
5 an elixir, or an emulsion.

A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active compound in a free flowing form, e.g., a powder or granules, optionally mixed with accessory ingredients, e.g., binders, lubricants, inert  
10 diluents, disintegrating agents or coloring agents. Molded tablets may be made by molding in a suitable machine a mixture of the active compound, preferably in powdered form, with a suitable carrier. Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethyl-  
15 cellulose, polyethylene glycol, waxes and the like. Lubricants used in these dosage forms include, without limitation, sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

20 Oral liquid forms, such as syrups or suspensions in suitably flavored suspending or dispersing agents such as the synthetic and natural gums, for example, tragacanth, acacia, methyl cellulose and the like, may be made by adding the active compound to the solution or suspension. Additional dispersing agents which may be employed include glycerin and the like.

25 Formulations for vaginal or rectal administration may be presented as a suppository with a conventional carrier, i.e., a base that is nontoxic and nonirritating to mucous membranes, compatible with a compound of structural formula I, and is stable in storage and does not bind or interfere with the release of the compound of structural formula I. Suitable bases include: cocoa butter (theobroma oil),  
30 polyethylene glycols (such as carbowax and polyglycols), glycol-surfactant

combinations, polyoxyl 40 stearate, polyoxyethylene sorbitan fatty acid esters (such as Tween, Myrj, and Arlacel), glycerinated gelatin, and hydrogenated vegetable oils. When glycerinated gelatin suppositories are used, a preservative such as methylparaben or propylparaben may be employed.

5                   Topical preparations containing the active drug component can be admixed with a variety of carrier materials well known in the art, such as, e.g., alcohols, aloe vera gel, allantoin, glycerine, vitamin A and E oils, mineral oil, PPG2 myristyl propionate, and the like, to form, e.g., alcoholic solutions, topical cleansers, cleansing creams, skin gels, skin lotions, and shampoos in cream or gel formulations.

10                   The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

                  Compounds of the present invention may also be delivered by the use  
15 of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxy-ethylaspartamidephenol, or polyethylene-oxide polylysine substituted  
20 with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

25                   Formulations suitable for parenteral administration include formulations that comprise a sterile aqueous preparation of the active compound which is preferably isotonic with the blood of the recipient. Such formulations suitably comprise a solution or suspension of a compound that is isotonic with the blood of the recipient subject. Such formulations may contain distilled water, 5%  
30 dextrose in distilled water or saline and the active compound. Often it is useful to

employ a pharmaceutically and pharmacologically acceptable acid addition salt of the active compound that has appropriate solubility for the solvents employed. Useful formulations also comprise concentrated solutions or solids comprising the active compound which on dilution with an appropriate solvent give a solution suitable for  
5 parenteral administration.

The compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates, and cross-linked or amphipathic  
10 block copolymers of hydrogels.

The pharmaceutical composition and method of the present invention may further comprise other therapeutically active compounds usually applied in the treatment of the above mentioned conditions, including osteoporosis, periodontal disease, bone fracture, bone damage following bone reconstructive surgery,  
15 sarcopenia, frailty, aging skin, male hypogonadism, post-menopausal symptoms in women, atherosclerosis, hypercholesterolemia, hyperlipidemia, aplastic anemia and other hematopoietic disorders, pancreatic cancer, Alzheimer's disease, inflammatory arthritis, and joint repair.

For the treatment and prevention of osteoporosis, the compounds of the present invention may be administered in combination with a bone-strengthening agent selected from antiresorptive agents, osteoanabolic agents, and other agents beneficial for the skeleton through mechanisms which are not precisely defined, such as calcium supplements, flavonoids, and vitamin D analogs. The conditions of periodontal disease, bone fracture, and bone damage following bone reconstructive  
20 surgery may also benefit from these combined treatments. For example, the compounds of the instant invention may be effectively administered in combination with effective amounts of other agents such as estrogens, bisphosphonates, SERMs, eathepsin K inhibitors,  $\alpha v \beta 3$  integrin receptor antagonists, vacuolar ATPase inhibitors, the polypeptide osteoprotegerin, antagonists of VEGF, thiazolidinediones,  
25 calcitonin, protein kinase inhibitors, parathyroid hormone (PTH) and analogs, calcium  
30

receptor antagonists, growth hormone secretagogues, growth hormone releasing hormone, insulin-like growth factor, bone morphogenetic protein (BMP), inhibitors of BMP antagonism, prostaglandin derivatives, fibroblast growth factors, vitamin D and derivatives thereof, vitamin K and derivatives thereof, soy isoflavones, calcium salts, and fluoride salts. The conditions of periodontal disease, bone fracture, and bone damage following bone reconstructive surgery may also benefit from these combined treatments. In one embodiment of the present invention, a compound of the instant invention may be effectively administered in combination with an effective amount of a bone-strengthening agent selected from the group consisting of estrogen or an estrogen derivative, alone or in combination with a progestin or progestin derivative; a bisphosphonate; an antiestrogen or a selective estrogen receptor modulator; an  $\alpha\beta3$  integrin receptor antagonist; a cathepsin K inhibitor; an osteoclast vacuolar ATPase inhibitor; calcitonin; and osteoprotegerin.

In the treatment of osteoporosis, the activity of the compounds of the present invention are distinct from that of the anti-resorptive agents: estrogens, bisphosphonates, SERMs, calcitonin, cathepsin K inhibitors, vacuolar ATPase inhibitors, agents interfering with the RANK/RANKL/Osteoprotegerin pathway, p38 inhibitors or any other inhibitors of osteoclast generation or osteoclast activation. Rather than inhibiting bone resorption, the compounds of structural formula I stimulate bone formation, acting preferentially on cortical bone, which is responsible for a significant part of bone strength. The thickening of cortical bone substantially contributes to a reduction in fracture risk, especially fractures of the hip. The combination of the tissue-selective androgen receptor modulators of structural formula I with anti-resorptive agents such as estrogen, bisphosphonates, antiestrogens, SERMs, calcitonin,  $\alpha\beta3$  integrin receptor antagonists, HMG-CoA reductase inhibitors, vacuolar ATPase inhibitors, and cathepsin K inhibitors is particularly useful because of the complementarity of the bone anabolic and antiresorptive actions.

Bone antiresorptive agents are those agents which are known in the art to inhibit the resorption of bone and include, for example, estrogen and estrogen derivatives which include steroidal compounds having estrogenic activity such as, for

example, 17 $\beta$ -estradiol, estrone, conjugated estrogen (PREMARIN®), equine estrogen, 17 $\beta$ -ethynyl estradiol, and the like. The estrogen or estrogen derivative may be employed alone or in combination with a progestin or progestin derivative. Nonlimiting examples of progestin derivatives are norethindrone and medroxy-  
5 progesterone acetate.

Bisphosphonates are also bone anti-resorptive agents. Bisphosphonate compounds which may also be employed in combination with a compound of structural formula I of the present invention include:

- (a) alendronic acid: (4-amino-1-hydroxybutylidene)-bis-phosphonic acid;
- 10 (b) alendronate (also known as alendronate sodium or monosodium trihydrate):  
(4-amino-1-hydroxybutylidene)-bis-phosphonate monosodium trihydrate (alendronic acid and alendronate are described in U.S. Patents 4,922,007, to Kieczkowski et al., issued May 1, 1990, and 5,019,651, to Kieczkowski, issued May 28, 1991, both of which  
15 are incorporated by reference herein in their entirety);
- (c) [(cycloheptylamino)-methylene]-bis-phosphonate (incadronate), which is described in U.S. Patent 4,970,335, to Isomura et al., issued November 13, 1990, which is incorporated by reference herein in I  
ts entirety;
- 20 (d) (dichloromethylene)-bis-phosphonic acid (clodronic acid) and the disodium salt (clodronate), which are described in Belgium Patent 672,205 (1966) and *J. Org. Chem* 32, 4111 (1967), both of which are incorporated by reference herein in their entirety;
- (e) [1-hydroxy-3-(1-pyrrolidinyl)-propylidene]-bis-phosphonate (EB-1053);
- 25 (f) (1-hydroxyethylidene)-bis-phosphonate (etidronate);
- (g) [1-hydroxy-3-(methylpentylamino)propylidene]-bis-phosphonate (ibandronate), which is described in U.S. Patent No. 4,927,814, issued May 22, 1990, which is incorporated by reference herein in I  
ts entirety;
- 30 (h) (6-amino-1-hydroxyhexylidene)-bis-phosphonate (neridronate);

- (i) [3-(dimethylamino)-1-hydroxypropylidene]-bis-phosphonate (olpadronate);  
(j) (3-amino-1-hydroxypropylidene)-bis-phosphonate (pamidronate);  
(k) [2-(2-pyridinyl)ethylidene]-bis-phosphonate (piridronate), which is described  
in U.S. Patent No. 4,761,406, which is incorporated by reference in  
5 its entirety;  
(l) [1-hydroxy-2-(3-pyridinyl)-ethylidene]-bis-phosphonate (risedronate);  
(m) {[4-(4-chlorophenyl)thio]methylene}-bis-phosphonate (tiludronate), which is  
described in U.S. Patent 4,876,248, to Breliere et al., October 24,  
1989, which is incorporated by reference herein in its entirety;  
10 (n) [1-hydroxy-2-(1H-imidazol-1-yl)ethylidene]-bis-phosphonate (zoledronate);  
and  
(o) [1-hydroxy-2-imidazopyridin-(1,2-a)-3-ylethylidene]-bis-phosphonate  
(minodronate).

In one embodiment of the methods and compositions of the present  
15 invention, the bisphosphonate is selected from the group consisting of alendronate,  
clodronate, etidronate, ibandronate, incadronate, minodronate, neridronate,  
olpadronate, pamidronate, piridronate, risedronate, tiludronate, and zoledronate, and  
pharmaceutically acceptable salts thereof, and mixtures thereof. In a class of this  
embodiment, the bisphosphonate is selected from the group consisting of alendronate,  
20 risedronate, zoledronate, ibandronate, tiludronate, and clodronate. In a subclass of  
this class, the bisphosphonate is alendronate, pharmaceutically acceptable salts and  
hydrates thereof, and mixtures thereof. A particular pharmaceutically acceptable salt  
of alendronate is alendronate monosodium. Pharmaceutically acceptable hydrates of  
alendronate monosodium include the monohydrate and the trihydrate. A particular  
25 pharmaceutically acceptable salt of risedronate is risedronate monosodium.  
Pharmaceutically acceptable hydrates of risedronate monosodium include the hemi-  
pentahydrate.

As used throughout this specification and claims, the term "alendronic  
acid" includes the related bisphosphonic acid forms, pharmaceutically acceptable salt  
30 forms, and equilibrium mixtures of these. It includes crystalline, hydrated crystalline,

and amorphous forms of alendronic acid and pharmaceutically acceptable salts thereof. It specifically includes anhydrous alendronate monosodium, alendronate monosodium monohydrate, and alendronate monosodium trihydrate.

Still further, antiestrogenic compounds such as raloxifene (see, e.g.,  
5 U.S. Pat. No. 5,393,763), clomiphene, zuclomiphene, enclomiphene, nafoxidene, CI-680, CI-628, CN-55,945-27, Mer-25, U-11,555A, U-100A, and salts thereof, and the like (see, e.g., U.S. Patent Nos. 4,729,999 and 4,894,373) may be employed in combination with a compound of structural formula I in the methods and compositions of the present invention. These agents are also known as SERMs, or  
10 selective estrogen receptor modulators, agents known in the art to prevent bone loss by inhibiting bone resorption via pathways believed to be similar to those of estrogens. These agents may be used in combination with the compounds of the present invention to beneficially treat bone disorders including osteoporosis. Such agents include, for example, tamoxifen, raloxifene, lasofoxifene, toremifene,  
15 azorxifene, EM-800, EM-652, TSE 424, clomiphene, droloxifene, idoxifene, and levormeloxifene [Goldstein, et al., "A pharmacological review of selective estrogen receptor modulators," Human Reproduction Update, 6: 212-224 (2000), and Lufkin, et al., "The role of selective estrogen receptor modulators in the prevention and treatment of osteoporosis," Rheumatic Disease Clinics of North America, 27: 163-185  
20 (2001)]. SERMs are also discussed in "Targeting the Estrogen Receptor with SERMs," Ann. Rep. Med. Chem. 36: 149-158 (2001).

$\alpha v \beta 3$  Integrin receptor antagonists suppress bone resorption and may be employed in combination with the tissue selective androgen receptor modulators of structural formula I for the treatment of bone disorders including osteoporosis.  
25 Peptidyl as well as peptidomimetic antagonists of the  $\alpha v \beta 3$  integrin receptor have been described both in the scientific and patent literature. For example, reference is made to W.J. Hoekstra and B.L. Poulter, Curr. Med. Chem. 5: 195-204 (1998) and references cited therein; WO 95/32710; WO 95/37655; WO 97/01540; WO 97/37655; WO 98/08840; WO 98/18460; WO 98/18461; WO 98/25892; WO 98/31359; WO  
30 98/30542; WO 99/15506; WO 99/15507; WO 00/03973; EP 853084; EP 854140; EP

- 854145; US Patent Nos. 5,204,350; 5,217,994; 5,639,754; 5,741,796; 5,780,426; 5,929,120; 5,952,341; 6,017,925; and 6,048,861. Evidence of the ability of  $\alpha v \beta 3$  integrin receptor antagonists to prevent bone resorption in vitro and in vivo has been presented (see V.W. Engleman et al., "A Peptidomimetic Antagonist of the  $\alpha v \beta 3$  Integrin Inhibits Bone Resorption In Vitro and Prevents Osteoporosis In Vivo," J. Clin. Invest. 99: 2284-2292 (1997); S.B. Rodan et al., "A High Affinity Non-Peptide  $\alpha v \beta 3$  Ligand Inhibits Osteoclast Activity In Vitro and In Vivo," J. Bone Miner. Res. 11: S289 (1996); J.F. Gourvest et al., "Prevention of OVX-Induced Bone Loss With a Non-peptidic Ligand of the  $\alpha v \beta 3$  Vitronectin Receptor," Bone 23: S612 (1998); M.W. Lark et al., "An Orally Active Vitronectin Receptor  $\alpha v \beta 3$  Antagonist Prevents Bone Resorption In Vitro and In Vivo in the Ovariectomized Rat," Bone 23: S219 (1998)). Other  $\alpha v \beta 3$  antagonists are described in R.M. Keenan et al., "Discovery of Potent Nonpeptide Vitronectin Receptor ( $\alpha v \beta 3$ ) Antagonists," J. Med. Chem. 40: 2289-2292 (1997); R.M. Keenan et al., "Benzimidazole Derivatives As Arginine Mimetics in 1,4-Benzodiazepine Nonpeptide Vitronectin Receptor ( $\alpha v \beta 3$ ) Antagonists," Bioorg. Med. Chem. Lett. 8: 3165-3170 (1998); and R.M. Keenan et al., "Discovery of an Imidazopyridine-Containing 1,4-Benzodiazepine Nonpeptide Vitronectin Receptor ( $\alpha v \beta 3$ ) Antagonist With Efficacy in a Restenosis Model," Bioorg. Med. Chem. Lett. 8: 3171-3176 (1998). Still other benzazepine, benzodiazepine and benzocycloheptene  $\alpha v \beta 3$  integrin receptor antagonists are described in the following patent publications: WO 96/00574, WO 96/00730, WO 96/06087, WO 96/26190, WO 97/24119, WO 97/24122, WO 97/24124, WO 98/14192, WO 98/15278, WO 99/05107, WO 99/06049, WO 99/15170, WO 99/15178, WO 99/15506, and U.S. Patent No. 6,159,964, and WO 97/34865.  $\alpha v \beta 3$  integrin receptor antagonists having dibenzocycloheptene, dibenzocycloheptane and dibenzoxazepine scaffolds have been described in WO 97/01540, WO 98/30542, WO 99/11626, WO 99/15508, WO 00/33838, U.S. Patent Nos. 6,008,213, and 6,069,158. Other osteoclast integrin receptor antagonists incorporating backbone conformational ring constraints have been described in the patent literature. Published patent applications or issued patents disclosing antagonists having a phenyl constraint include WO 98/00395, WO

99/32457, WO 99/37621, WO 99/44994, WO 99/45927, WO 99/52872, WO 99/52879, WO 99/52896, WO 00/06169, EP 0 820,988, EP 0 820,991, U.S. Patent Nos. 5,741,796; 5,773,644; 5,773,646; 5,843,906; 5,852,210; 5,929,120; 5,952,381; 6,028,223; and 6,040,311. Published patent applications or issued patents disclosing antagonists having a monocyclic ring constraint include WO 99/26945, WO 99/30709, WO 99/30713, WO 99/31099, WO 99/59992, WO 00/00486, WO 00/09503, EP 0 796,855, EP 0 928,790, EP 0 928,793, U.S. Patent Nos. 5,710,159; 5,723,480; 5,981,546; 6,017,926; and 6,066,648. Published patent applications or issued patents disclosing antagonists having a bicyclic ring constraint include WO 98/23608, WO 98/35949, WO 99/33798, EP 0 853,084, U.S. Patent Nos. 5,760,028; 5,919,792; and 5,925,655. Reference is also made to the following reviews for additional scientific and patent literature that concern  $\alpha_v$  integrin antagonists: M. E. Duggan, et al., "Ligands to the integrin receptor  $\alpha_v\beta_3$ , Exp. Opin. Ther. Patents, 10: 1367-1383 (2000); M. Gowen, et al., "Emerging therapies for osteoporosis," Emerging Drugs, 5: 1-43 (2000); J.S. Kerr, et al., "Small molecule  $\alpha_v$  integrin antagonists: novel anticancer agents," Exp. Opin. Invest. Drugs, 9: 1271-1291 (2000); and W.H. Miller, et al., "Identification and in vivo efficacy of small-molecule antagonists of integrin  $\alpha_v\beta_3$  (the vitronectin receptor)," Drug Discovery Today, 5: 397-408 (2000).

20                   Cathepsin K, formerly known as cathepsin O2, is a cysteine protease and is described in PCT International Application Publication No. WO 96/13523, published May 9, 1996; U.S. Patent No. 5,501,969, issued March 3, 1996; and U.S. Patent No. 5,736,357, issued April 7, 1998, all of which are incorporated by reference herein in their entirety. Cysteine proteases, specifically cathepsins, are linked to a number of disease conditions, such as tumor metastasis, inflammation, arthritis, and bone remodeling. At acidic pH's, cathepsins can degrade type-I collagen. Cathepsin protease inhibitors can inhibit osteoclastic bone resorption by inhibiting the degradation of collagen fibers and are thus useful in the treatment of bone resorption diseases, such as osteoporosis.

Members of the class of HMG-CoA reductase inhibitors, known as the "statins," have been found to trigger the growth of new bone, replacing bone mass lost as a result of osteoporosis (see The Wall Street Journal, Friday, December 3, 1999, page B1). Therefore, the statins hold promise for the treatment of bone resorption.

- 5 Examples of HMG-CoA reductase inhibitors include statins in their lactonized or dihydroxy open acid forms and pharmaceutically acceptable salts and esters thereof, including but not limited to lovastatin (see US Patent No. 4,342,767); simvastatin (see US Patent No. 4,444,784); dihydroxy open-acid simvastatin, particularly the ammonium or calcium salts thereof; pravastatin, particularly the sodium salt thereof  
10 (see US Patent No. 4,346,227); fluvastatin, particularly the sodium salt thereof (see US Patent No. 5,354,772); atorvastatin, particularly the calcium salt thereof (see US Patent No. 5,273,995); cerivastatin, particularly the sodium salt thereof (see US Patent No. 5,177,080), rosuvastatin, also known as ZD-4522 (see US Patent No. 5,260,440) and pitavastatin, also referred to as NK-104, itavastatin, or nisvastatin (see PCT  
15 international application publication number WO 97/23200).

Osteoclast vacuolar ATPase inhibitors, also called proton pump inhibitors, may also be employed together with the tissue selective androgen receptor modulators of structural formula I. The proton ATPase which is found on the apical membrane of the osteoclast has been reported to play a significant role in the bone  
20 resorption process. Therefore, this proton pump represents an attractive target for the design of inhibitors of bone resorption which are potentially useful for the treatment and prevention of osteoporosis and related metabolic diseases [see C. Farina et al., "Selective inhibitors of the osteoclast vacuolar proton ATPase as novel bone antiresorptive agents," DDT, 4: 163-172 (1999)].

25 The angiogenic factor VEGF has been shown to stimulate the bone-resorbing activity of isolated mature rabbit osteoclasts via binding to its receptors on osteoclasts [see M. Nakagawa et al., "Vascular endothelial growth factor (VEGF) directly enhances osteoclastic bone resorption and survival of mature osteoclasts," FEBS Letters, 473: 161-164 (2000)]. Therefore, the development of antagonists of

VEGF binding to osteoclast receptors, such as KDR/Flk-1 and Flt-1, may provide yet a further approach to the treatment or prevention of bone resorption.

Activators of the peroxisome proliferator-activated receptor- $\gamma$ (PPAR $\gamma$ ), such as the thiazolidinediones (TZD's), inhibit osteoclast-like cell formation and bone resorption *in vitro*. Results reported by R. Okazaki *et al.* in Endocrinology, 140: 5060-5065 (1999) point to a local mechanism on bone marrow cells as well as a systemic one on glucose metabolism. Nonlimiting examples of PPAR $\gamma$  activators include the glitazones, such as troglitazone, pioglitazone, rosiglitazone, and BRL 49653.

Calcitonin may also be employed together with the tissue selective androgen receptor modulator of structural formula I. Calcitonin is preferentially employed as salmon nasal spray (Azra *et al.*, Calcitonin. 1996. In: J. P. Bilezikian, *et al.*, Ed., Principles of Bone Biology, San Diego: Academic Press; and Silverman, "Calcitonin," Rheumatic Disease Clinics of North America, 27: 187-196, 2001)

Protein kinase inhibitors may also be employed together with the tissue selective androgen receptor modulators of structural formula I. Kinase inhibitors include those disclosed in WO 01/17562 and are in one embodiment selected from inhibitors of p38. Specific embodiments of p38 inhibitors useful in the present invention include SB 203580 [Badger *et al.*, "Pharmacological profile of SB 203580, a selective inhibitor of cytokine suppressive binding protein/p38 kinase, in animal models of arthritis, bone resorption, endotoxin shock, and immune function," J. Pharmacol. Exp. Ther., 279: 1453-1461 (1996)].

Osteoanabolic agents are those agents that are known in the art to build bone by increasing the production of the bone protein matrix. Such osteoanabolic agents include, for example, the various forms of parathyroid hormone (PTH) such as naturally occurring PTH (1-84), PTH (1-34), analogs thereof, native or with substitutions and particularly parathyroid hormone subcutaneous injection. PTH has been found to increase the activity of osteoblasts, the cells that form bone, thereby promoting the synthesis of new bone (Modern Drug Discovery, Vol. 3, No. 8, 2000). In studies reported at the First World Congress on Osteoporosis held in Chicago in

June 2000, women in combined PTH-estrogen therapy exhibited a 12.8% increase in spinal bone mass and a 4.4% increase in total hip mass. Another study presented at the same meeting showed that PTH could increase bone size as well as density. A clinical trial of the effect of the human parathyroid hormone 1-34 fragment [hPTH(1-34)] on postmenopausal osteoporotic women resulted in  $\geq 65\%$  reduction in spine fractures and a 54% reduction in nonvertebral fractures, after a median of 22 months of treatment [see J.M. Hock, Bone, 27: 467-469 (2000) and S. Mohan, et al., Bone, 27: 471-478 (2000), and references cited therein]. Thus, PTH and fragments thereof, such as hPTH(1-34), may prove to be efficacious in the treatment of osteoporosis alone or in combination with other agents, such as the tissue selective androgen receptor modulators of the present invention. An injectable recombinant form of human PTH, Forteo (teriparatide), has received regulatory approval in the U.S. for the treatment of osteoporosis.

Also useful in combination with the SARMs of the present invention are calcium receptor antagonists which induce the secretion of PTH as described by Gowen et al., in "Antagonizing the parathyroid calcium receptor stimulates parathyroid hormone secretion and bone formation in osteopenic rats," J. Clin. Invest. 105: 1595-604 (2000).

Growth hormone secretagogues, growth hormone, growth hormone releasing hormone and the like are also osteoanabolic agents which may be employed with the compounds according to structural formula I for the treatment of osteoporosis. Representative growth hormone secretagogues are disclosed in U.S. Patent No. 3,239,345; U.S. Patent No. 4,036,979; U.S. Patent No. 4,411,890; U.S. Patent No. 5,206,235; U.S. Patent No. 5,283,241; U.S. Patent No. 5,284,841; U.S. Patent No. 5,310,737; U.S. Patent No. 5,317,017; U.S. Patent No. 5,374,721; U.S. Patent No. 5,430,144; U.S. Patent No. 5,434,261; U.S. Patent No. 5,438,136; U.S. Patent No. 5,494,919; U.S. Patent No. 5,494,920; U.S. Patent No. 5,492,916; U.S. Patent No. 5,536,716; EPO Patent Pub. No. 0,144,230; EPO Patent Pub. No. 0,513,974; PCT Patent Pub. No. WO 94/07486; PCT Patent Pub. No. WO 94/08583; PCT Patent Pub. No. WO 94/11012; PCT Patent Pub. No. WO 94/13696; PCT Patent

Pub. No. WO 94/19367; PCT Patent Pub. No. WO 95/03289; PCT Patent Pub. No. WO 95/03290; PCT Patent Pub. No. WO 95/09633; PCT Patent Pub. No. WO 95/11029; PCT Patent Pub. No. WO 95/12598; PCT Patent Pub. No. WO 95/13069; PCT Patent Pub. No. WO 95/14666; PCT Patent Pub. No. WO 95/16675; PCT Patent Pub. No. WO 95/16692; PCT Patent Pub. No. WO 95/17422; PCT Patent Pub. No. WO 95/17423; PCT Patent Pub. No. WO 95/34311; PCT Patent Pub. No. WO 96/02530; Science, 260, 1640-1643 (June 11, 1993); Ann. Rep. Med. Chem., 28: 177-186 (1993); Bioorg. Med. Chem. Lett., 4: 2709-2714 (1994); and Proc. Natl. Acad. Sci. USA, 92: 7001-7005 (1995).

10                   Insulin-like growth factor (IGF) may also be employed together with the tissue selective androgen receptor modulators of structural formula I. Insulin-like growth factors may be selected from Insulin-like Growth Factor I, alone or in combination with IGF binding protein 3 and IGF II [See Johannson and Rosen, "The IGFs as potential therapy for metabolic bone diseases," 1996, In: Bilezikian, et al.,  
15   Ed., Principles of Bone Biology, San Diego: Academic Press; and Ghiron et al., "Effects of recombinant insulin-like growth factor-I and growth hormone on bone turnover in elderly women," J. Bone Miner. Res. 10: 1844-1852 (1995)].

                  Bone morphogenetic protein (BMP) may also be employed together with the tissue selective androgen receptor modulators of structural formula I. Bone  
20   morphogenetic protein includes BMP 2, 3, 5, 6, 7, as well as related molecules TGF beta and GDF 5 [Rosen et al., "Bone morphogenetic proteins," 1996. In: J. P. Bilezikian, et al., Ed., Principles of Bone Biology, San Diego: Academic Press; and Wang EA, "Bone morphogenetic proteins (BMPs): therapeutic potential in healing bony defects," Trends Biotechnol., 11: 379-383 (1993)].

25                   Inhibitors of BMP antagonism may also be employed together with the tissue selective androgen receptor modulators of structural formula I. BMP antagonist inhibitors are in one embodiment selected from inhibitors of the BMP antagonists SOST, noggin, chordin, gremlin, and dan [Massague and Chen, "Controlling TGF-beta signaling," Genes Dev., 14: 627-644, 2000; Aspenberg et al., "The bone  
30   morphogenetic proteins antagonist Noggin inhibits membranous ossification," J. Bone

Miner. Res. 16: 497-500, 2001; Brunkow et al., "Bone dysplasia sclerosteosis results from loss of the SOST gene product, a novel cystine knot-containing protein," Am. J. Hum. Genet. 68: 577-89 (2001)].

5 The tissue-selective androgen receptor modulators of the present invention may also be combined with the polypeptide osteoprotegerin for the treatment of conditions associated with bone loss, such as osteoporosis. Preferably osteoprotegerin is mammalian osteoprotegerin and more preferably human osteoprotegerin. The polypeptide osteoprotegerin, a member of the tumor necrosis factor receptor superfamily, is useful to treat bone diseases characterized by increased  
10 bone loss, such as osteoporosis. Reference is made to U.S. Patent No. 6,288,032, which is incorporated by reference herein in its entirety.

Prostaglandin derivatives may also be employed together with the tissue selective androgen receptor modulators of structural formula I. Prostaglandin derivatives are in one embodiment selected from agonists of prostaglandin receptor  
15 EP1, EP2, EP4, FP and IP or a derivative thereof [Pilbeam et al., "Prostaglandins and bone metabolism," 1996. In: Bilezikian, et al. Ed. Principles of Bone Biology, San Diego: Academic Press; Weinreb et al., "Expression of the prostaglandin E(2) (PGE(2)) receptor subtype EP(4) and its regulation by PGE(2) in osteoblastic cell lines and adult rat bone tissue," Bone, 28: 275-281 (2001)].

20 Fibroblast growth factors may also be employed together with the tissue selective androgen receptor modulators of structural formula I. Fibroblast growth factors include aFGF, bFGF and related peptides with FGF activity [Hurley Florkiewicz, "Fibroblast growth factor and vascular endothelial growth factor families," 1996. In: J. P. Bilezikian, et al., Ed. Principles of Bone Biology, San Diego:  
25 Academic Press].

In addition to bone resorption inhibitors and osteoanabolic agents, there are also other agents known to be beneficial for the skeleton through mechanisms which are not precisely defined. These agents may also be favorably combined with the tissue selective androgen receptor modulators of structural formula  
30 I.

Vitamin D and vitamin D derivatives may also be employed together with the tissue selective androgen receptor modulator of structural formula I. Vitamin D and vitamin D derivatives include natural vitamin D, 25-OH-vitamin D3, 1 $\alpha$ ,25(OH)<sub>2</sub> vitamin D3, 1 $\alpha$ -OH-vitamin D3, 1 $\alpha$ -OH-vitamin D2, dihydrotachysterol, 26,27-F6-1 $\alpha$ ,25(OH)<sub>2</sub> vitamin D3, 19-nor-1 $\alpha$ ,25(OH)<sub>2</sub> vitamin D3, 22-oxacalcitriol, calcipotriol, 1 $\alpha$ ,25(OH)<sub>2</sub>-16-ene-23-yne-vitamin D3 (Ro 23-7553), EB1089, 20-epi-1 $\alpha$ ,25(OH)<sub>2</sub> vitamin D3, KH1060, ED71, 1 $\alpha$ ,24(S)-(OH)<sub>2</sub> vitamin D3, 1 $\alpha$ ,24(R)-(OH)<sub>2</sub> vitamin D3 [See, Jones G., "Pharmacological mechanisms of therapeutics: vitamin D and analogs," 1996. In: J. P. Bilezikian, et al. Ed. Principles of Bone Biology, San Diego: Academic Press].

Vitamin K and vitamin K derivatives may also be employed together with the tissue selective androgen receptor modulators of structural formula I. Vitamin K and vitamin K derivatives include menatetrenone (vitamin K2) [see Shiraki et al., "Vitamin K2 (menatetrenone) effectively prevents fractures and sustains lumbar bone mineral density in osteoporosis," J. Bone Miner. Res., 15: 515-521 (2000)].

Soy isoflavones, including ipriflavone, may be employed together with the tissue selective androgen receptor modulators of structural formula I.

Fluoride salts, including sodium fluoride (NaF) and monosodium fluorophosphate (MFP), may also be employed together with the tissue selective androgen receptor modulators of structural formula I. Dietary calcium supplements may also be employed together with the tissue selective androgen receptor modulators of structural formula I. Dietary calcium supplements include calcium carbonate, calcium citrate, and natural calcium salts (Heaney. Calcium. 1996. In: J. P. Bilezikian, et al., Ed., Principles of Bone Biology, San Diego: Academic Press).

Daily dosage ranges for bone resorption inhibitors, osteoanabolic agents and other agents which may be used to benefit the skeleton when used in combination with a compound of structural formula I are those which are known in the art. In such combinations, generally the daily dosage range for the tissue selective androgen receptor modulator of structural formula I is 0.01 to 1000 mg per adult

human per day, more preferably from 0.1 to 200 mg/day. However, adjustments to decrease the dose of each agent may be made due to the increased efficacy of the combined agent.

5 In particular, when a bisphosphonate is employed, dosages of 2.5 to 100 mg/day (measured as the free bisphosphonic acid) are appropriate for treatment, more preferably 5 to 20 mg/day, especially about 10 mg/day. Prophylactically, doses of about 2.5 to about 10 mg/day and especially about 5 mg/day should be employed. For reduction in side-effects, it may be desirable to administer the combination of a compound of structural formula I and the bisphosphonate once a week. For once  
10 weekly administration, doses of about 15 mg to 700 mg per week of bisphosphonate and 0.07 to 7000 mg of a compound of structural formula I may be employed, either separately, or in a combined dosage form. A compound of structural formula I may be favorably administered in a controlled-release delivery device, particularly for once weekly administration.

15 For the treatment of atherosclerosis, hypercholesterolemia, and hyperlipidemia, the compounds of structural formula I may be effectively administered in combination with one or more additional active agents. The additional active agent or agents can be lipid-altering compounds such as HMG-CoA reductase inhibitors, or agents having other pharmaceutical activities, or agents that  
20 have both lipid-altering effects and other pharmaceutical activities. Examples of HMG-CoA reductase inhibitors include statins in their lactonized or dihydroxy open acid forms and pharmaceutically acceptable salts and esters thereof, including but not limited to lovastatin (see US Patent No. 4,342,767); simvastatin (see US Patent No. 4,444,784); dihydroxy open-acid simvastatin, particularly the ammonium or calcium  
25 salts thereof; pravastatin, particularly the sodium salt thereof (see US Patent No. 4,346,227); fluvastatin, particularly the sodium salt thereof (see US Patent No. 5,354,772); atorvastatin, particularly the calcium salt thereof (see US Patent No. 5,273,995); cerivastatin, particularly the sodium salt thereof (see US Patent No. 5,177,080), and nisvastatin, also referred to as NK-104 (see PCT international  
30 application publication number WO 97/23200). Additional active agents which may

be employed in combination with a compound of structural formula I include, but are not limited to, HMG-CoA synthase inhibitors; squalene epoxidase inhibitors; squalene synthetase inhibitors (also known as squalene synthase inhibitors), acyl-coenzyme A: cholesterol acyltransferase (ACAT) inhibitors including selective inhibitors of ACAT-1 or ACAT-2 as well as dual inhibitors of ACAT-1 and -2; microsomal triglyceride transfer protein (MTP) inhibitors; probucol; niacin; cholesterol absorption inhibitors, such as SCH-58235, also known as ezetimibe and 1-(4-fluorophenyl)-3(R)-[3(S)-(4-fluorophenyl)-3-hydroxypropyl]-4(S)-(4-hydroxyphenyl)-2-azetidinone, which is described in U.S. Patent Nos. 5,767,115 and 5,846,966; bile acid sequestrants; LDL (low density lipoprotein) receptor inducers; platelet aggregation inhibitors, for example glycoprotein IIb/IIIa fibrinogen receptor antagonists and aspirin; human peroxisome proliferator activated receptor gamma (PPAR $\gamma$ ) agonists, including the compounds commonly referred to as glitazones, for example troglitazone, pioglitazone and rosiglitazone and, including those compounds included within the structural class known as thiazolidinediones as well as those PPAR $\gamma$  agonists outside the thiazolidinedione structural class; PPAR $\alpha$  agonists, such as clofibrate, fenofibrate including micronized fenofibrate, and gemfibrozil; PPAR dual  $\alpha/\gamma$  agonists; vitamin B<sub>6</sub> (also known as pyridoxine) and the pharmaceutically acceptable salts thereof such as the HCl salt; vitamin B<sub>12</sub> (also known as cyanocobalamin); folic acid or a pharmaceutically acceptable salt or ester thereof such as the sodium salt and the methylglucamine salt; anti-oxidant vitamins such as vitamin C and E and beta carotene; beta-blockers; angiotensin II antagonists such as losartan; angiotensin converting enzyme inhibitors, such as enalapril and captopril; calcium channel blockers, such as nifedipine and diltiazem; endothelin antagonists; agents such as LXR ligands that enhance ABC1 gene expression; bisphosphonate compounds, such as alendronate sodium; and cyclooxygenase-2 inhibitors, such as rofecoxib and celecoxib, as well as other agents known to be useful in the treatment of these conditions.

Daily dosage ranges for HMG-CoA reductase inhibitors when used in combination with the compounds of structural formula I correspond to those which

are known in the art. Similarly, daily dosage ranges for the HMG-CoA synthase inhibitors; squalene epoxidase inhibitors; squalene synthetase inhibitors (also known as squalene synthase inhibitors), acyl-coenzyme A: cholesterol acyltransferase (ACAT) inhibitors including selective inhibitors of ACAT-1 or ACAT-2 as well as  
5 dual inhibitors of ACAT-1 and -2; microsomal triglyceride transfer protein (MTP) inhibitors; probucol; niacin; cholesterol absorption inhibitors including ezetimibe; bile acid sequestrants; LDL (low density lipoprotein) receptor inducers; platelet aggregation inhibitors, including glycoprotein IIb/IIIa fibrinogen receptor antagonists and aspirin; human peroxisome proliferator activated receptor gamma (PPAR $\gamma$ )  
10 agonists; PPAR $\alpha$  agonists; PPAR dual  $\alpha/\gamma$  agonists; vitamin B<sub>6</sub>; vitamin B<sub>12</sub>; folic acid; anti-oxidant vitamins; beta-blockers; angiotensin II antagonists; angiotensin converting enzyme inhibitors; calcium channel blockers; endothelin antagonists; agents such as LXR ligands that enhance ABC1 gene expression; bisphosphonate compounds; and cyclooxygenase-2 inhibitors also correspond to those which are  
15 known in the art, although due to the combined action with the compounds of structural formula I, the dosage may be somewhat lower when administered in combination.

In accordance with the method of the present invention, the individual components of the combination can be administered separately at different times  
20 during the course of therapy or concurrently in divided or single combination forms. The instant invention is therefore to be understood as embracing all such regimes of simultaneous or alternating treatment and the term "administering" is to be interpreted accordingly. It will be understood that the scope of combinations of the compounds of this invention with other agents useful for treating diseases caused by androgen  
25 deficiency or that can be ameliorated by addition of androgen.

Abbreviations Used in the Description of the Preparation of the Compounds of the Present Invention:

AcOH	Acetic acid
AgOTf	Silver triflate
calc.	Calculated
CH <sub>2</sub> Cl <sub>2</sub>	Methylene chloride
DEAD	Diethyl azodicarboxylate
DIEA	Diisopropylethylamine
DMAP	4-Dimethylaminopyridine
DMF	N,N-Dimethylformamide
DMSO	Dimethyl sulfoxide
EDC.HCl	1-(3-Dimethylaminopropyl)3-ethylcarbodiimide HCl
ES-MS	Electron-spray mass spectroscopy
Et	ethyl
Et <sub>2</sub> O	Diethyl ether
Et <sub>3</sub> N	Triethylamine
EtOAc	Ethyl acetate
FAB	Fast atom bombardment
HOBt	N-hydroxybenzotriazole
HPLC	High-performance liquid chromatography
HRMS	High resolution mass spectrum
i-PrOH	Isopropyl alcohol
K <sub>2</sub> CO <sub>3</sub>	Potassium carbonate
Me	Methyl
Me <sub>2</sub> SO <sub>4</sub>	Dimethyl sulfate
MF	molecular formula
MgSO <sub>4</sub>	Magnesium sulfate
MS	mass spectrum
NBS	N-bromosuccinimide

NMM	N-methylmorpholine
Na <sub>2</sub> SO <sub>4</sub>	Sodium sulfate
Pd(PPh <sub>3</sub> ) <sub>4</sub>	Tetrakis(triphenylphosphine)palladium(0)
Ph	Phenyl
Pr	Propyl
rt	Room temperature
NaHCO <sub>3</sub>	Sodium hydrogencarbonate
PPh <sub>3</sub>	Triphenylphosphine
THF	Tetrahydrofuran
TLC	Thin-layer chromatography

Preparation of the Compounds of the Invention

The compounds of the present invention can be prepared according to the procedures denoted in the following reaction Schemes and Examples or modifications thereof using readily available starting materials, reagents, and



5 conventional procedures or variations thereof well-known to a practitioner of ordinary skill in the art of synthetic organic chemistry. Specific definitions of variables in the Schemes are given for illustrative purposes only and are not intended to limit the procedures described.

The following examples are provided to further illustrate details for the


10 preparation and use of the compounds of the present invention. They are not intended to be limitations on the scope of the instant invention in any way, and they should not be so construed. Furthermore, the compounds described in the following examples are not to be construed as forming the only genus that is considered as the invention, and any combination of the compounds or their moieties may itself form a genus.

15 Those skilled in the art will readily understand that known variations of the conditions and processes of the following preparative procedures can be used to prepare these compounds. All temperatures are in degrees Celsius unless noted otherwise.

## EXAMPLE 1



5



Step A:      3-Oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxylic acid methyl ester (1-2)

A mixture of 1-1, (*J. Med. Chem.*, 29: 2298-2315 (1986)), (50.0 g, 157.5 mmol), EDC·HCl (33.2 g, 173.3 mmol) and DMAP (1.9 g, 15.8 mmol) in MeOH (300 mL) was stirred for 24 hr. The mixture was concentrated and diluted  
5 with water (1000 mL). After filtration, the solid was collected and dried to furnish the desired product 1-2 as a solid which was used in Step B without further purification.  
MS calculated M+H: 332.2, found 332.2

Step B:      4-Methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxylic acid methyl ester (1-3)

To a suspension of 1-2 (7.0 g, 21.1 mmol) in 100 mL dry THF was added NaH (1.3 g, 32 mmol) gradually. The reaction mixture was stirred at rt for 1 h. Me<sub>2</sub>SO<sub>4</sub> (10 mL) was added in one portion. The mixture was stirred overnight.  
MeOH (30 mL) was gradually added. After stirring for 3 h, water (500 mL) was  
15 added. A solid precipitated out immediately. After filtration, the collected solid was dissolved in chloroform and washed with brine and dried (MgSO<sub>4</sub>). After solvent removal, the desired product 1-3 was collected as a solid and used in Step C without further purification.  
MS calculated M+H: 346.2, found 346.3

20

Step C:      6-Bromo-4-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxylic acid methyl ester (1-4)

To a suspension of 1-3 (11 g, 33 mmol) in CCl<sub>4</sub> (100 mL) was added NBS (6.2 g, 45 mmol) and benzoyl peroxide (0.1 g, 0.5 mmol). The reaction mixture  
25 was refluxed for 4 h. The suspension was cooled and filtered. The filtrate was concentrated to provide the desired product 1-4, which was used in Step D without further purification.  
MS calculated M+H: 424.1, found 424.2

Step D:        4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxylic acid  
                         methyl ester (1-5)

---


A mixture of 1-4 (0.8 g, 1.8 mmol), K<sub>2</sub>CO<sub>3</sub> (0.7 g, 5.3 mmol), Pd(PPh<sub>3</sub>)<sub>4</sub> (0.2 g, 0.2 mmol) and trimethylboroxine (0.3 mL, 2.1 mmol) in DMF (15 mL) was purged with nitrogen for 10 min. The mixture was then heated at 100 °C overnight. The reaction was quenched with saturated aqueous NaHCO<sub>3</sub> and then water (200 mL). It was extracted three times with EtOAc. The combined organic layers were washed with water, brine and dried (MgSO<sub>4</sub>). After solvent removal, the residue was purified by silica gel flash chromatography (100% EtOAc to 80% EtOAc/20% Hexanes) to afford the desired product 1-5 as a solid.  
MS calculated M+H: 360.3, found 360.3

Step E:        4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxylic  
                         acid (1-6)

---

To a solution of 1-5 (1.9 g, 5.3 mmol) in 1,4-dioxane (20 mL) was added NaOH (0.5 g) in 5 mL water. The mixture was refluxed overnight and acidified with 3N HCl and diluted with 50 mL water. The suspension was extracted three times with chloroform. The combined organic layers were washed with brine and dried (MgSO<sub>4</sub>) and concentrated to provide the desired product 1-6 as a solid which was used in Step F with further purification.  
MS calculated M+H: 347.2, found 346.3

## EXAMPLE 2



S-(Pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carbothioate (2-1)

A mixture of 1-6 (1.9 g, 5.3 mmol), Ph<sub>3</sub>P (2.8 g, 10.6 mmol) and 2,2'-dipyridyl disulfide (2.3 g, 10.6 mmol) in toluene (30 mL) was stirred overnight. After solvent removal, the residue was purified by flash silica gel chromatography (10% EtOAc/90% Hexanes to 100% EtOAc) to provide the desired product 2-1 as a solid. MS calculated M+H: 439.2, found 439.3


N-Phenyl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide (2-2)

To a solution of 2-1 (0.05 g, 0.1 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1 mL) was added aniline (0.03 g, 0.3 mmol) and AgOTf (0.3 g, 0.1 mmol). The mixture was stirred










overnight. After filtration, the filtrate was concentrated and subsequently purified via reverse-phase chromatography [30% CH<sub>3</sub>CN/70% H<sub>2</sub>O (with 0.02% HCl) to 95% CH<sub>3</sub>CN/5% H<sub>2</sub>O (with 0.02% HCl) over 10 min] to afford the desired product 2-2.  
MS calculated M+H: 421.3, found 421.3








- 5                    Examples 2-84 in Table 1 were prepared in a similar manner as Compound 2-2, (Example 2 in Table 1), but using the appropriate R<sup>2</sup>R<sup>3</sup>NH amine to generate the final desired carboxamide.









TABLE 1




















EX	NR <sup>2</sup> R <sup>3</sup>	NAME	Mass spectrum Measured [M + 1]
2		N-phenyl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	421.3
3		N-(2-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	455.3
4		N-(3-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	455.3
5		N-(4-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	455.3








6		N-(2-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	489.3
7		N-(3-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	489.3
8		N-(4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	489.3
9		N-(2-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	451.3
10		N-(3-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	451.3
11		N-(4-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	451.3
12		N-(2,2,2-trifluoroethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	427.3
13		N-(2-fluorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	453.3
14		N-(3-fluorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	453.3







15		N-(4-fluorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	453.3
16		N-(pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	436.3
17		N-(2,3-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	525.3
18		N-(2,5-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	525.2530
19		N-(2-chloro-6-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	523.2318
20		N-(2-chloro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	523.2314
21		N-(4-chloro-2-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	523.2315








22		N-(2,4-dichloro-5-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	519.2176
23		N-[4-(morpholin-4-yl)phenyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	506.3379
24		N-(3-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	505.2676
25		N-(4-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	505.2660
26		N-(2-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	505.3
27		N-[2-(piperidin-1-yl)phenyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	504.3574
28		N-(5-fluoro-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	440.2698
29		N-(2-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	439.274









30		N-(4-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	435.2997
31		N-(5-methyl-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	436.2954
32		N-(4-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	439.2741
33		N-(2-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	435.2992
34		N-(3-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	435.2994
35		N-(3-cyanophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	446.2797
36		N-(pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	422.2806
37		N-(pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	422.2807
38		N-(pyridin-3-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	436.2952









39		N-(2-chloro-4-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	473.2355
40		N-(2,6-dichlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	489.2067
41		N-(2,6-difluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	457.2658
42		N-(2,3,4,6-tetrafluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	493.2463
43		N-(2,3-dichlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	489.2063
44		N-(2,5-dichlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	489.2064
45		N-(thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	428.2366
46		N-(4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	446.2790



47		N-(3-chloro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	480.2397
48		N-(3-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	480.2402
49		N-(3-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	480.2398
50		N-(4-acetyl-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	463.2944
51		N-(4-methyl-thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	442.2502
52		N-(4-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	480.2393
53		N-(2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	446.2782

54		N-(2,5-difluoro-4-cyano-phenyl)- 4-methyl-6-methyl-3-oxo-4-aza- 5α-androst-5-en-17β-carboxamide	482.2589
55		N-(2,4,5,6-tetrafluoro-4-cyano- phenyl)-4-methyl-6-methyl-3-oxo- 4-aza-5α-androst-5-en-17β- carboxamide	518.2394
56		N-(2,6-dichloro-4-cyano-phenyl)- 4-methyl-6-methyl-3-oxo-4-aza- 5α-androst-5-en-17β-carboxamide	514.2018
57		N-(3,4-dicyano-phenyl)-4-methyl- 6-methyl-3-oxo-4-aza-5α-androst- 5-en-17β-carboxamide	471.2743
58		N-(2,4-dichloro-6-cyano-phenyl)- 4-methyl-6-methyl-3-oxo-4-aza- 5α-androst-5-en-17β-carboxamide	514.2016
59		N-(2-chloro-4-cyano-phenyl)-4- methyl-6-methyl-3-oxo-4-aza-5α- androst-5-en-17β-carboxamide	480.2391

60		N-(3-fluoro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	464.2696
61		N-(3-trifluoromethyl-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	514.1
62		N-(2-chloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	456.2403
63		N-(3,5-dichloro-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	490.2019
64		N-(2-chloro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	456.2405
65		N-(2,5-dichloro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	490.2017
66		N-(3-chloro-5-trifluoromethyl-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	524.2268


67		N-(2,6-dichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	490.2027
68		3-[[[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino]-4-chlorobenzoic acid	499.2360
69		N-(4-chlorobenzothiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	512.2135
70		N-(benzothiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	478.2527
71		N-(4,5-dimethylthiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	456.2678
72		N-(2,3,5-trichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	524.1639
73		N-(2,6-dichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	490.2025
74		N-(5-ethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	457.2625

75		N-(4-methyl-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	436.1
76		N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	469.2617
77		N-(2-chloro-6-trifluoromethyl-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	524.2257
78		N-(2,6-dichloro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	490.2004
79		N-(2,6-dichloro-6'-trifluoro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	558.2
80		N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	443.2458
81		N-(4-methyl-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	436.2997
82		N-(pyrazin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	423.2762

83		N-(6-chloro-pyrazin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	457.2334
84		N-(2-methyl-3-chloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	470.0

## EXAMPLE 85

5



N-(1H-benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide (3-1)

10


A mixture of 1-6 (0.15 g, 0.4 mmol), RNH<sub>2</sub> (0.26 g, 1.3 mmol), EDC (0.17 g, 0.87 mmol), HOAt (0.089 g, 0.65 mmol) and EtN<sup>i</sup>Pr<sub>2</sub> (0.28 g, 2.2 mmol) was stirred at rt overnight. It was concentrated and purified via reverse phase HPLC to afford the desired product 3-1 as a solid. MS calculated M+H: 475.652, found

15

475.3054







Examples 86-130 in Table 2 were prepared in a similar manner as Compound 3-1 (Example 85 in Table 2), but using the appropriate amine to generate the final desired carboxamide.






TABLE 2













5







EX	NR <sup>2</sup> R <sup>3</sup>	NAME	MASS SPECTRUM Measured [M + 1]
85		N-(benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	475.3054
86		N-(2-fluoro-6-chlorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	487.0
87		N-((R)-phenylmethylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	449.1







88		N-((S)-phenylmethylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	449.1
89		N-(2,6-dichlorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	503.2229
90		N-(3,4-dichlorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	503.2222
91		N-[(1S,2S)-2-hydroxycyclohexyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	443.3252
92		N-(3H-imidazo[4,5-b]pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	476.3000
93		N-[(1R,2S)-2-hydroxycyclohexyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	443.3251








94		N-(1,3-thiazol-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	428.2368
95		N-(1H-indol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	474.3106
96		N-[2-(1H-benzimidazol-2-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	489.3204
97		N-[2-(6-chloro-1H-benzimidazol-2-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	523.2809
98		tert-butyl 3-({[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino}methyl)-1H-indole-1-carboxylate	574.0




99		tert-butyl 3-(2- {[(1,4a,6a,11-tetramethyl- 2-oxo- 2,3,4,4a,4b,5,6,6a,7,8,9,9a ,9b,10-tetradecahydro-1H- indeno[5,4-f]quinolin-7- yl)carbonyl]amino}ethyl)- 1H-indole-1-carboxylate	588.0
100		N-(benzimidazol-2- ylmethylmethyl)-4- methyl-6-methyl-3-oxo-4- aza-5α-androst-5-en-17β- carboxamide	489.3201
101		N-[2-(1-methyl-1H- benzimidazol-2-yl)ethyl]- 4-methyl-6-methyl-3-oxo- 4-aza-5α-androst-5-en- 17β-carboxamide	503.3359
102		N-(imidazo[1,2- a]pyrimidin-2-ylmethyl)- 4-methyl-6-methyl-3-oxo- 4-aza-5α-androst-5-en- 17β-carboxamide	476.3010
103		N-(1H-indol-5-ylmethyl)- 4-methyl-6-methyl-3-oxo- 4-aza-5α-androst-5-en- 17β-carboxamide	474.3107

104		N-[(6-chloro-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	509.2665
105		N-(imidazo[1,2-a]pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	475.3056
106		N-(2-(imidazo[1,2-a]pyridin-2-ylethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	489.3208
107		N-[2-(2-phenyl-1H-indol-3-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	564.3567
108		N-(1-methyl-6-trifluoromethyl-1H-benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	557.3067


109		N-(1H-indol-7-ylmethyl)- 4-methyl-6-methyl-3-oxo- 4-aza-5 $\alpha$ -androst-5-en- 17 $\beta$ -carboxamide	474.3103
110		N-(1H-indol-4-ylmethyl)- 4-methyl-6-methyl-3-oxo- 4-aza-5 $\alpha$ -androst-5-en- 17 $\beta$ -carboxamide	474.3101
111		N-(imidazo[1,2-a]pyridin- 2-ylmethyl)-4-methyl-6- methyl-3-oxo-4-aza-5 $\alpha$ - androst-5-en-17 $\beta$ - carboxamide	475.3062
112		N-[(1-phenyl-1H- benzimidazol-2- yl)methyl]-4-methyl-6- methyl-3-oxo-4-aza-5 $\alpha$ - androst-5-en-17 $\beta$ - carboxamide	551.3363
113		N-(pyrazolo[1,5- a]pyridin-7-ylmethyl)-4- methyl-6-methyl-3-oxo-4- aza-5 $\alpha$ -androst-5-en-17 $\beta$ - carboxamide	475.3059
114		N-(pyrazolo[1,5- a]pyridin-2-ylmethyl)-4- methyl-6-methyl-3-oxo-4- aza-5 $\alpha$ -androst-5-en-17 $\beta$ - carboxamide	475.3061

115		N-[(1-methyl-2,3-dihydro-1H-indol-5-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	490.3414
116		N-piperidin-4-yl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	428.3258
117		N-piperidin-3-yl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	428.3289
118		N-(2-ethylpyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	450.1
119		N-[(1-methyl-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	489.3208
120		N-(5-tert-Butyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	485.0

121		N-(5-trifluoromethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	497.0
122		N-(2-methoxypyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	452.0
123		N-(2-ethoxypyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	466.0
124		N-(pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	422.2790
125		N-(4-trifluoromethylpyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	490.2643
126		N-(4,6-dimethylpyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	450.3116
127		N-(6-methylpyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-	436.2932

		17 $\beta$ -carboxamide	
128		N-(7H-purin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	463.2775
129		N-(5-cyclopropyl-1H-pyrazol-3-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	465.3206
130		N-(6-methoxybenzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide	505.3155

## EXAMPLE 131



5

Step A: Methyl (3S,3aS,6R)-6-(3-methoxy-3-oxopropyl)-3a,6-dimethyl-7-oxododecahydro-1H-cyclopenta[a]naphthalene-3-carboxylate (4-1)

To a suspension of 1-1 (50 g, 157.5 mmol) in 400 mL methanol cooled at 0 °C was added conc  $\text{H}_2\text{SO}_4$  (50 mL) slowly. After the addition, the reaction mixture was heated to 50 °C overnight. After removal of the solvent, the residue was poured into ice. The mixture was extracted with EtOAc (x3). The combined organic layers was washed with water, sat  $\text{NaHCO}_3$ , brine and dried ( $\text{MgSO}_4$ ). Removal of the solvent afforded a yellow oil which was dissolved in 100 mL methanol. The solution was then cooled to 0 °C. A cooled (0 °C) solution of

NaOH (6.5 g, 159.7 mmol) was added. The mixture was stirred overnight, treated with 3N HCl and concentrated. The residue was extracted with CH<sub>2</sub>Cl<sub>2</sub> (x3). The combined organic layers was washed with water, brine and dried (MgSO<sub>4</sub>). Removal of the solvent afforded the desired product 4-1 as a white solid, which was used for the next step without further purification.

MS calculated M+H: 351.5, found 351.0

Step B: 3-[(3S,3aS,6R)-3-(methoxycarbonyl)-3a,6-dimethyl-8-methylene-7-oxododecahydro-1H-cyclopenta[a]naphthalen-6-yl]propanoic acid (4-2)

To a solution of 4-1 ( 24.5 g, 69.9 mmol) in 250 mL dichloromethane was added the Eschenmoser's salt (16.4 g, 174.8 mmol). The mixture was refluxed over 48 hr and then cooled to room temperature. After addition of MeI ( 8.7 mL, 139.8 mmol) and Et<sub>3</sub>N (24.4 mL, 174.8 mmol), the mixture was stirred at room temperature for 48 hr and concentrated. The residue was dissolved in dichloromethane (150 mL). DBU (21 mL, 139.8 mmol) was added. The mixture was stirred overnight and diluted with water (150 mL) and acidified with 6N HCl until pH=4. After removal of the solvent, the residue was extracted with EtOAc (x3). The combined organic layers was washed with brine and dried (MgSO<sub>4</sub>). Concentration of the mixture afforded a yellow solid as the desired product 4-2 which was further purified by recrystallization (methanol/water).

MS calculated M-H<sub>2</sub>O: 344.5, found 345.0

Step C: 3-[(3S,3aS,6R)-8,8-ethylene-3-(methoxycarbonyl)-3a,6-dimethyl-7-oxododecahydro-1H-cyclopenta[a]naphthalen-6-yl]propanoic acid (4-3).

To a suspension of trimethylsulfoxonium bromide (35.8 g, 206.9 mmol) in 180 mL DMSO was added NaH (5.0 g, 206.9 mmol) gradually (bubbles). The mixture was stirred for 1.5 hr. A solution of 4-2 (18.7 g, 51.7 mmol) in 70 mL DMSO was added rapidly via cannula and stirred for 1 hr. The mixture was poured

into ice and extracted with EtOAc (x3). After removal of solvent, the residue was purified on reverse phase HPLC with gradient (5% CH<sub>3</sub>CN/75% H<sub>2</sub>O (0.02%)/20% MeOH) to (75% CH<sub>3</sub>CN/5% H<sub>2</sub>O (0.02%)/20% MeOH) over 10 min. Upon concentration, the desired product 4-3 was obtained a beige solid.

5 MS calculated M+1: 377.5, found 377.0

Step D: 6,6-Ethylene-5-hydroxy-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxylic acid methyl ester (4-4)

To a solution of 4-3 (10.0 g, 26.6 mmol) in 100 mL dichloromethane at 0 °C was added Et<sub>3</sub>N (14.8 mL, 106.2 mmol) and trimethylacetyl chloride (3.9 mL, 31.9 mmol) and catalytic The mixture was stirred for 1 hr. To the reaction mixture was added a solution of anhydrous ammonia in dichloromethane (prepared by bubbling anhydrous ammonia into dichloromethane for 50 min at 0 °C). The mixture was stirred for 2 hr at room temperature and followed by treatment with aqueous

10 at 0 °C was added Et<sub>3</sub>N (14.8 mL, 106.2 mmol) and trimethylacetyl chloride (3.9 mL, 31.9 mmol) and catalytic The mixture was stirred for 1 hr. To the reaction mixture was added a solution of anhydrous ammonia in dichloromethane (prepared by bubbling anhydrous ammonia into dichloromethane for 50 min at 0 °C). The mixture was stirred for 2 hr at room temperature and followed by treatment with aqueous

15 NaHCO<sub>3</sub>. The organic layer was separated and dried (MgSO<sub>4</sub>). Upon removal of the solvent. The desired product 4-4 was obtained as a beige solid which was used for the next step without further purification.

MS calculated M+1: 376.5, found 376.1

20 Step E: 6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxylic acid methyl ester (4-5)

To a solution of 4-4 (6.4 g, 17.0 mmol) in 55 mL dichloromethane was added Et<sub>3</sub>SiH (110 mL) and 55 mL TFA. The mixture was stirred at rt for 2 h and concentrated. The residue was treated with sat. NaHCO<sub>3</sub> and extracted with EtOAc (x3). The combined organic layers was washed with sat. NaHCO<sub>3</sub>, brine and dried (MgSO<sub>4</sub>). After removal of the solvent, the desired product 4-5 was obtained as a colorless oil, which was used for the next step without further purification.

25 (x3). The combined organic layers was washed with sat. NaHCO<sub>3</sub>, brine and dried (MgSO<sub>4</sub>). After removal of the solvent, the desired product 4-5 was obtained as a colorless oil, which was used for the next step without further purification.

MS calculated M+1: 360.5, found 360.1

Step F:            6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxylic acid (4-6)

- To a suspension of 4-5 (4.6 g, 12.8 mmol) in 80 mL methanol was added a solution of LiOH (1.6 g, 38.4 mL). The mixture was heated at 50 °C overnight. The reaction was quenched with 3N HCl until pH =3. The mixture was extracted with a mixed solvent (chloroform/isopropanol) three times. The combined organic layers was dried (MgSO<sub>4</sub>). Upon removal of the solvent, the desired product 4-6 was obtained as a beige solid, which was used in the next step without further purification.
- 10    MS calculated M+1: 346.5, found 346.0.

Step G:            S-(Pyridin-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carbothioate(4-7)


- A mixture of 4-6 (0.5 g, 1.3 mmol), Aldrithiol-2 (0.6 g, 2.6 mmol) and triphenylphosphine (0.7 g, 2.6 mmol) in 5 mL toluene was stirred overnight. After removal of the solvent, the desired product 4-7 and other side products were obtained as a yellow syrup. Due to the difficulty of separation, the desired product was used in the next step without further purification.
- 15    MS calculated M+1: 439.6, LC/MS found 439.2

Step H:            N-(2',3'-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide (4-8)

- To a solution of 4-7 (0.6 g crude material prepared above) was added silver triflate (1.7 g, 6.4 mmol) and 2, 3-difluoroaniline (0.8 g, 6.4 mmol). The mixture was stirred at rt overnight and filtered. The solution was concentrated and purified via reverse phase HPLC with gradient (5% CH<sub>3</sub>CN/75% H<sub>2</sub>O (0.02%)/20% MeOH) to (75% CH<sub>3</sub>CN/5% H<sub>2</sub>O (0.02%)/20% MeOH) over 10 min to afford the desired product 4-8 as a solid.
- 25    MS calculated M+1: 457.6, LC/MS found 457.0






Examples 132-156 in Table 3 were prepared in a similar manner as Compound 4-8 (Example 131 in Table 3) using the appropriate amine to generate the final desired carboxamide. Some of these examples were prepared using the reaction conditions similar to those in Example 84 to couple compound 4-6 with the  
5 appropriate amine.






TABLE 3













5




EX	NHR	NAME	Mass spectrum Measured [M + 1]
131		N-(2,3-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	457.2655
132		N-(2-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	455.2447
133		N-(2,6-dichlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	490.2

134		N-(2-fluorophenyl)- 6,6-Ethylene-3-oxo- 4-aza-5 $\alpha$ -androst- 17 $\beta$ -carboxamide	439.2748
135		N-(2-fluoro-6- chlorophenyl)-6,6- Ethylene-3-oxo-4- aza-5 $\alpha$ -androst- 17 $\beta$ -carboxamide	473.2360
136		N-(2-fluoro-6- fluorophenyl)-6,6- Ethylene-3-oxo-4- aza-5 $\alpha$ -androst- 17 $\beta$ -carboxamide	457.2656
137		N-(2- trifluoromethyl-6- chlorophenyl)-6,6- Ethylene-3-oxo-4- aza-5 $\alpha$ -androst- 17 $\beta$ -carboxamide	523.2330
138		N-(2-fluoro-4- chlorophenyl)-6,6- Ethylene-3-oxo-4- aza-5 $\alpha$ -androst- 17 $\beta$ -carboxamide	473.2367


139		N-(2-trifluoromethylphenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	489.2705
140		N-(2-fluoro-5-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	473.2368
141		N-phenyl-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	421.3
142		N-(5-ethyl-1,3,4-thiadiazol-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	457.2601
143		N-(5-methyl-1,3,4-thiadiazol-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	443.2497

144		N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	469.2627
145		N-(pyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	422.2795
146		N-(2-chloropyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	456.2396
147		N-(2-methylpyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	436.2944
148		N-(2-chloropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	456.2408

149		N-(4-fluoropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	440.2696
150		N-(4-chloropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	456.2413
151		N-(4-methylpyridin-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	436.2923
152		N-(4-chloropyridin-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	456.2413
153		N-(benzimidazol-2-ylmethyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	475.3081

154		N-((N-methylbenzimidazol-1-yl)methyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	489.3209
155		N-(3H-imidazo[4,5-b]pyridin-2-yl)methyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	476.3046
156		N-(pyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide	422.2811

## EXAMPLE 157



Step A:      6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxylic acid  
methyl ester (5-1)

---

- To a solution of 4-5 (0.05 g, 0.14 mmol) in 5 mL THF was added
- 5    NaOtBu in one portion. The mixture was stirred for 1 hr at rt. Me<sub>2</sub>SO<sub>4</sub> (0.05 mL, 0.6 mmol) was added neat. The mixture was stirred overnight and treated with water. The solution was extracted with ethyl ether (x3). The combined organic layers was washed with brine and dried. After concentration, the desired product 5-1 was obtained a solid which was used in the next step without further purification.
- 10   MS calculated M+1: 374.0, LC/MS found 374.0

Step B:      6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -  
carboxylic acid (5-2)

---

- To a solution of 5-1 (1.8 g, 4.8 mmol) in 7 mL methanol was added
- 15    LiOH (1.2 g, 42.9 mmol) in 5 mL water. The reaction was heated at 50 °C for 4 hr, cooled to rt and quenched with 6N HCl. The mixture was extracted with CHCl<sub>3</sub> (x3). The combined organic layers was washed with brine and dried (MgSO<sub>4</sub>). After removal of the solvent, the desired product 6-2 was obtained as a solid.
- MS calculated M+1: 360.5, LC/MS found 360.3

20

Step C:      S-(Pyridin-2-yl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-  
17 $\beta$ -carbothioate (5-3)

---

- A mixture of 5-2 (1.4 g, 4.0 mmol), Aldrithiol-2 (1.8 g, 8.0 mmol) and triphenylphosphine (2.1 g, 8.0 mmol) in 15 mL toluene was stirred at rt overnight.
- 25    After removal of the solvent, the desired product 5-3 and other products were obtained as a mixture, which was used in the next step without further purification due to the difficulty of separation.
- MS calculated M+1: 453.66, LC/MS found 453.0


Step D:      N-(2',3'-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -  
androst-17 $\beta$ -carboxamide (5-4)

---

- To a solution of 5-3 (0.6 g crude material prepared above) was added  
5 silver triflate (1.7 g, 6.6 mmol) and 2, 3-difluoroaniline (0.9 g, 6.6 mmol). The  
mixture was stirred at rt overnight and filtered. The solution was concentrated and  
purified via reverse phase HPLC with gradient (5% CH<sub>3</sub>CN/85%H<sub>2</sub>O  
(0.02%)/10%MeOH) to (85% CH<sub>3</sub>CN/5%H<sub>2</sub>O (0.02%)/10%MeOH) over 10 min to  
afford the desired product 5-4 as a solid.  
10 MS calculated M+1: 471.6, MS found 471.0





Examples 158-168 in Table 4 were prepared in a similar manner as  
Compound 5-4 (Example 157 in Table 4) but using the appropriate amine to generate  
the final desired carboxamide.






Table 4



5

EX	NHR	NAME	MASS SPECTRU M Measured [M + 1]
157		N-(2,3-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide	471.2812
158		N-(2-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide	469.2605
159		N-(2,6-dichlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -	503.2228

		carboxamide	
160		N-(2-fluorophenyl)- 6,6-Ethylene-3-oxo- 4-aza-4-methyl-5 $\alpha$ - androst-17 $\beta$ - carboxamide	453.2900
161		N-(2-fluoro-6- chlorophenyl)-6,6- Ethylene-3-oxo-4- aza-4-methyl-5 $\alpha$ - androst-17 $\beta$ - carboxamide	487.2508
162		N-(2,6- difluorophenyl)-6,6- Ethylene-3-oxo-4- aza-4-methyl-5 $\alpha$ - androst-17 $\beta$ - carboxamide	471.2807
163		N-(2-trifluoromethyl- 6-chlorophenyl)-6,6- Ethylene-3-oxo-4- aza-4-methyl-5 $\alpha$ - androst-17 $\beta$ - carboxamide	537.2478

164		N-(2-fluoro-4-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide	487.2506
165		N-(2-trifluoromethylphenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide	503.2865
166		N-(2-fluoro-5-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide	487.2518
167		N-phenyl-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide	435.3
168		N-(2,6-dichloro-4-cyanophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide	

## EXAMPLE 169

Oral Composition

- 5                   As a specific embodiment of an oral composition of a compound of this invention, 50 mg of a compound of the present invention is formatted with sufficient finely divided lactose to provide a total amount of 580 to 590 mg to fill a size 0 hard gelatin capsule.

10

## EXAMPLE 170

Transdermal Patch Formulation

<u>Ingredient</u>	<u>Amount</u>
Compound of formula I	40 g
Silicone fluid	45 g
Colloidal silicone dioxide	2.5 g

- The silicone fluid and compound of structural formula I are mixed together and the colloidal silicone dioxide is added to increase viscosity. The material is then dosed
- 15   into a subsequently heat sealed polymeric laminate comprised of the following: polyester release liner, skin contact adhesive composed of silicone or acrylic polymers, a control membrane which is a polyolefin (e.g. polyethylene, polyvinyl acetate or polyurethane), and an impermeable backing membrane made of a polyester multilaminate. The resulting laminated sheet is then cut into 10 cm<sup>2</sup> patches. For 100
- 20   Patches.

## EXAMPLE 171

5    Suppository

---

<u>Ingredient</u>	<u>Amount</u>
Compound of structural formula I	25 g
Polyethylene glycol 1000	1481 g
Polyethylene glycol 4000	494 g

The polyethylene glycol 1000 and polyethylene glycol 4000 are mixed and melted. The compound of structural formula I is mixed into the molten mixture, poured into molds and allowed to cool. For 1000 suppositories.

10

## EXAMPLE 172

Injectable solution

---

<u>Ingredient</u>	<u>Amount</u>
Compound of structural formula I	5 g
Buffering agents	q.s.
Propylene glycol	400 mg
Water for injection	600 mL

The compound of structural formula I and buffering agents are dissolved in the propylene glycol at about 50°C. The water for injection is then added with stirring

15    and the resulting solution is filtered, filled into ampules, sealed and sterilized by autoclaving. For 1000 Ampules.

## EXAMPLE 173

Injectable solution

<u>Ingredient</u>	<u>Amount</u>
Compound of structural formula I	5 g
Buffering agents	q.s.
Magnesium sulfate heptahydrate	100 mg
Water for injection	880 mL

- 5 The compound of structural formula I, magnesium sulfate heptahydrate and buffering agents are dissolved in the water for injection with stirring, and the resulting solution is filtered, filled into ampoules, sealed and sterilized by autoclaving. For 1000 Ampoules.

- 10 The following assays were used to characterize the activity of the tissue selective androgen receptor modulators of the present invention.

#### IN VITRO AND IN VIVO ASSAYS FOR IDENTIFICATION OF COMPOUNDS WITH SARM ACTIVITY

15

1. Hydroxylapatite-based Radioligand Displacement Assay of Compound Affinity for  
Endogenously Expressed AR

Materials:

*Binding Buffer:* TEGM (10 mM Tris-HCl, 1 mM EDTA, 10% glycerol, 1 mM beta-mecaptoethanol, 10 mM Sodium Molybdate, pH 7.2)

- 20 *50% HAP Slurry:* Calbiochem Hydroxylapatite, Fast Flow, in 10 mM Tris, pH 8.0 and 1 mM EDTA.

*Wash Buffer:* 40 mM Tris, pH7.5, 100 mM KCl, 1 mM EDTA and 1 mM EGTA.

95% EtOH

- 25 Methyltrienolone, [17 $\alpha$ -methyl-<sup>3</sup>H], (R1881\*); NEN NET590

Methyltrienolone (R1881), NEN NLP005 (dissolve in 95% EtOH)

Dihydrotestosterone (DHT) [1,2,4,5,6,7-<sup>3</sup>H(N)] NEN NET453

Hydroxylapatite Fast Flow; Calbiochem Cat#391947

Molybdate = Molybdic Acid (Sigma, M1651)

5

MDA-MB-453 cell culture media:

RPMI 1640 (Gibco 11835-055)

w/23.8 mM NaHCO<sub>3</sub>, 2 mM L-glutamine

In 500 mL of complete media      Final conc.

10      10 mL (1M Hepes)      20 mM

5 mL (200 mM L-glu)      4 mM

0.5 mL (10 mg/mL human insulin)      10 µg/mL

in 0.01 N HCl Calbiochem#407694-S)

50 mL FBS (Sigma F2442)      10%

15      1 mL (10 mg/mL Gentamicin      20 µg /mL

Gibco#15710-072)

Cell Passaging:

Cells (Hall R. E., et al., European Journal of Cancer, 30A: 484-490

20 (1994)) are rinsed twice in PBS, phenol red-free Trypsin-EDTA is diluted in the same  
PBS 1:10. The cell layers are rinsed with 1X Trypsin, extra Trypsin is poured out,  
and the cell layers are incubated at 37°C for ~ 2 min. The flask is tapped and checked  
for signs of cell detachment. Once the cells begin to slide off the flask, the complete  
media is added to kill the trypsin. The cells are counted at this point, then diluted to  
25 the appropriate concentration and split into flasks or dishes for further culturing  
(Usually 1:3 to 1:6 dilution).

Preparation of MDA-MB-453 Cell Lysate

When the MDA cells are 70 to 85% confluent, they are detached as  
described above, and collected by centrifuging at 1000 g for 10 min at 4°C. The cell  
30 pellet is washed twice with TEGM (10 mM Tris-HCl, 1 mM EDTA, 10% glycerol, 1

mM beta-mercaptoethanol, 10 mM Sodium Molybdate, pH 7.2). After the final wash, the cells are resuspended in TEGM at a concentration of  $10^7$  cells/mL. The cell suspension is snap frozen in liquid nitrogen or ethanol/dry ice bath and transferred to  $-80^{\circ}\text{C}$  freezer on dry ice. Before setting up the binding assay, the frozen samples are left on ice-water to just thaw (~1 hr). Then the samples are centrifuged at 12,500 g to 20,000 g for 30 min at  $4^{\circ}\text{C}$ . The supernatant is used to set-up assay right away. If using 50  $\mu\text{L}$  of supernatant, the test compound can be prepared in 50  $\mu\text{L}$  of the TEGM buffer.

Procedure for Multiple Compound Screening:

1x TEGM buffer is prepared, and the isotope-containing assay mixture is prepared in the following order: EtOH (2% final concentration in reaction),  $^3\text{H}$ -R1881 or  $^3\text{H}$ -DHT (0.5 nM final Conc. in reaction) and 1x TEGM. [eg. For 100 samples, 200  $\mu\text{L}$  (100 x 2) of EtOH + 4.25  $\mu\text{L}$  of 1:10  $^3\text{H}$ -R1881 stock + 2300  $\mu\text{L}$  (100 x 23) 1x TEGM]. The compound is serially diluted, e.g., if starting final conc. is 1  $\mu\text{M}$ , and the compound is in 25  $\mu\text{L}$  of solution, for duplicate samples, 75  $\mu\text{L}$  of 4x1  $\mu\text{M}$  solution is made and 3  $\mu\text{L}$  of 100  $\mu\text{M}$  is added to 72  $\mu\text{L}$  of buffer, and 1:5 serial dilution.

25 $\mu\text{L}$  of  $^3\text{H}$ -R1881 trace and 25  $\mu\text{L}$  compound solution are first mixed together, followed by addition of 50  $\mu\text{L}$  receptor solution. The reaction is gently mixed, spun briefly at about 200 rpm and incubated at  $4^{\circ}\text{C}$  overnight. 100  $\mu\text{L}$  of 50% HAP slurry is prepared and added to the incubated reaction which is then vortexed and incubated on ice for 5 to 10 minutes. The reaction mixture is vortexed twice more to resuspend HAP while incubating reaction. The samples in 96-well format are then washed in wash buffer using The FilterMate<sup>TM</sup> Universal Harvester plate washer (Packard). The washing process transfers HAP pellet containing ligand-bound expressed receptor to Unifilter-96 GF/B filter plate (Packard). The HAP pellet on the filter plate is incubated with 50  $\mu\text{L}$  of MICROSCINT (Packard) scintillant for 30 minutes before being counted on the TopCount microscintillation counter (Packard).

IC<sub>50</sub>s are calculated using R1881 as a reference. Tissue selective androgen receptor modulators of the present invention displayed IC<sub>50</sub> values of 1 micromolar or less.

## 2. MMP1 Promoter Suppression, Transient Transfection Assay (TRAMPS)

5                   HepG2 cells are cultured in phenol red free MEM containing 10 %  
charcoal-treated FCS at 37C with 5% CO<sub>2</sub>. For transfection, cells are plated at  
10,000 cells/well in 96 well white, clear bottom plates. Twenty four hours later, cells  
are co-transfected with a MMP1 promoter-luciferase reporter construct and a rhesus  
monkey expression construct (50 : 1 ratio) using FuGENE6 transfection reagent,  
10 following the protocol recommended by manufacturer. The MMP1 promoter-  
luciferase reporter construct is generated by insertion of a human MMP1 promoter  
fragment (-179/+63) into pGL2 luciferase reporter construct (Promega) and a rhesus  
monkey AR expression construct is generated in a CMV-Tag2B expression vector  
(Stratagene). Cells are further cultured for 24 hours and then treated with test  
15 compounds in the presence of 100 nM phorbol-12-myristate-13-acetate (PMA), used  
to increase the basal activity of MMP1 promoter. The compounds are added at this  
point, at a range of 1000nM to 0.03nM, 10 dilutions, at a concentration on 10X,  
1/10th volume (example: 10 microliters of ligand at 10X added to 100 microliters of  
media already in the well). Cells are further cultured for an additional 48 hours. Cells  
20 are then washed twice with PBS and lysed by adding 70 µL of Lysis Buffer (1x,  
Promega) to the wells. The luciferase activity is measured in a 96-well format using a  
1450 Microbeta Jet (Perkin Elmer) luminometer. Activity of test compounds is  
presented as suppression of luciferase signal from the PMA-stimulated control levels.  
EC<sub>50</sub> and Emax values are reported. Tissue selective androgen receptor modulators  
25 of the present invention activate repression typically with submicromolar EC<sub>50</sub> values  
and Emax values greater than about 50%.

### References:

a. Newberry EP, Willis D, Latifi T, Boudreaux JM, Towler DA, "Fibroblast  
30 growth factor receptor signaling activates the human interstitial collagenase

promoter via the bipartite Ets-AP1 element," Mol. Endocrinol. 11: 1129-44 (1997).

- b. Schneikert J, Peterziel H, Defossez PA, Klocker H, Launoit Y, Cato AC,  
"Androgen receptor-Ets protein interaction is a novel mechanism for steroid  
5 hormone-mediated down-modulation of matrix metalloproteinase expression,"  
J. Biol. Chem. 271: 23907-23913 (1996).

3. A Mammalian Two-Hybrid Assay for the Ligand-induced Interaction of N-  
Terminus and C-Terminus Domains of the Androgen Receptor (Agonist  
10 Mode)

This assay assesses the ability of AR agonists to induce the interaction  
between the N-terminal domain (NTD) and C-terminal domain (CTD) of rhAR that  
reflects the in vivo virilizing potential mediated by activated androgen receptors. The  
interaction of NTD and CTD of rhAR is quantified as ligand induced association  
15 between a Gal4DBD-rhARCTD fusion protein and a VP16-rhARNTD fusion protein  
as a mammalian two-hybrid assay in CV-1 monkey kidney cells.

The day before transfection, CV-1 cells are trypsinized and counted,  
and then plated at 20,000 cells/well in 96-well plates or larger plates (scaled up  
accordingly) in DMEM + 10% FCS. The next morning, CV-1 cells are cotransfected  
20 with pCBB1 (Gal4DBD-rhARLBD fusion construct expressed under the SV40 early  
promoter), pCBB2 (VP16 -rhAR NTD fusion construct expressed under the SV40  
early promoter) and pFR (Gal4 responsive luciferase reporter, Promega) using  
LIPOFECTAMINE PLUS reagent (GIBCO-BRL) following the procedure  
recommended by the vendor. Briefly, DNA admixture of 0.05 µg pCBB1, 0.05 µg  
25 pCBB2 and 0.1 µg of pFR is mixed in 3.4 µL OPTI-MEM (GIBCO-BRL) mixed with  
"PLUS Reagent" (1.6µ L, GIBCO-BRL) and incubated at room temperature (RT) for  
15 min to form the pre-complexed DNA.

For each well, 0.4 µL LIPOFECTAMINE Reagent (GIBCO-BRL) is  
diluted into 4.6 µL OPTI-MEM in a second tube and mixed to form the diluted  
30 LIPOFECTAMINE Reagent. The pre-complexed DNA (above) and the diluted

LIPOFECTAMINE Reagent (above) are combined, mixed and incubated for 15 min at RT. The medium on the cells is replaced with 40  $\mu$ L /well OPTI-MEM, and 10  $\mu$ L DNA-lipid complexes are added to each well. The complexes are mixed into the medium gently and incubated at 37°C at 5% CO<sub>2</sub> for 5h. Following incubation, 200 L  
5  $\mu$ /well D-MEM and 13% charcoal-stripped FCS are added, followed by incubation at 37°C at 5% CO<sub>2</sub>. After 24 hours, the test compounds are added at the desired concentration(s) (1 nM – 10  $\mu$ M). Forty eight hours later, luciferase activity is measured using LUC-Screen system (TROPIX) following the manufacturer's protocol. The assay is conducted directly in the wells by sequential addition of 50  $\mu$ L  
10 each of assay solution 1 followed by assay solution 2. After incubation for 40 minutes at room temperature, luminescence is directly measured with 2-5 second integration.

Activity of test compounds is calculated as the E<sub>max</sub> relative to the activity obtained with 3 nM R1881. Typical tissue-selective androgen receptor modulators of the present invention display weak or no agonist activity in this assay  
15 with less than 50% agonist activity at 10 micromolar.

#### Reference:

He B, Kempainen JA, Voegel JJ, Gronemeyer H, Wilson EM, "Activation function in the human androgen receptor ligand binding domain mediates inter-domain  
20 communication with the NH(2)-terminal domain," J. Biol. Chem. 274: 37219-37225 (1999).

#### 4. A Mammalian Two-Hybrid Assay For Inhibition of Interaction between N-Terminus and C-Terminus Domains of Androgen Receptor (Antagonist 25 Mode)

This assay assesses the ability of test compounds to antagonize the stimulatory effects of R1881 on the interaction between NTD and CTD of rhAR in a mammalian two-hybrid assay in CV-1 cells as described above.

Forty eight hours after transfection, CV-1 cells are treated with test  
30 compounds, typically at 10  $\mu$ M, 3.3  $\mu$ M, 1  $\mu$ M, 0.33  $\mu$ M, 100  $\mu$ M, 33 nM, 10 nM, 3.3

nM and 1 nM final concentrations. After incubation at 37°C at 5% CO<sub>2</sub> for 10 - 30 minutes, an AR agonist methyltrienolone (R1881) is added to a final concentration of 0.3 nM and incubated at 37°C. Forty-eight hours later, luciferase activity is measured using LUC-Screen system (TROPIX) following the protocol recommended by the manufacturer. The ability of test compounds to antagonize the action of R1881 is calculated as the relative luminescence compared to the value with 0.3 nM R1881 alone.

SARM compounds of the present invention typically displayed antagonist activity in the present assay with IC<sub>50</sub> values less than 1 micromolar.

10

#### 5. Trans-Activation Modulation of Androgen Receptor (TAMAR)

This assay assesses the ability of test compounds to control transcription from the MMTV-LUC reporter gene in MDA-MB-453 cells, a human breast cancer cell line that naturally expresses the human AR. The assay measures induction of a modified MMTV LTR/promoter linked to the LUC reporter gene.

20,000 to 30,000 cells/well are plated in a white, clear-bottom 96-well plate in "Exponential Growth Medium" which consists of phenol red-free RPMI 1640 containing 10%FBS, 4mM L-glutamine, 20mM HEPES, 10ug/mL human insulin, and 20ug/mL gentamicin. Incubator conditions are 37°C and 5% CO<sub>2</sub>. The transfection is done in batch mode. The cells are trypsinized and counted to the right cell number in the proper amount of fresh media, and then gently mixed with the Fugene/DNA cocktail mix and plated onto the 96-well plate. All the wells receive 200 µl of medium + lipid/DNA complex and are then incubated at 37°C overnight. The transfection cocktail consists of serum-free Optimem, Fugene6 reagent and DNA.

The manufacturer's (Roche Biochemical) protocol for cocktail setup is followed. The lipid (µl) to DNA (µg) ratio is approximately 3:2 and the incubation time is 20 min at room temperature. Sixteen to 24 hrs after transfection, the cells are treated with test compounds such that the final DMSO (vehicle) concentration is <3%. The cells are exposed to the test compounds for 48 hrs. After 48 hrs, the cells are lysed by a

Promega cell culture lysis buffer for 30-60 min and then the luciferase activity in the extracts is assayed in the 96-well format luminometer.

Activity of test compounds is calculated as the  $E_{\max}$  relative to the activity obtained with 100 nM R1881.

5

References:

- a. R.E. Hall, et al., "MDA-MB-453, an androgen-responsive human breast carcinoma cell line with high androgen receptor expression," Eur. J. Cancer, 30A: 484-490 (1994).
- 10 b. R.E. Hall, et al., "Regulation of androgen receptor gene expression by steroids and retinoic acid in human breast-cancer cells," Int. J. Cancer, 52: 778-784 (1992).

6. In Vivo Prostate Assay

Male Sprague-Dawley rats aged 9-10 weeks, the earliest age of sexual maturity, are used in prevention mode. The goal is to measure the degree to which androgen-like compounds delay the rapid deterioration (~85%) of the ventral prostate gland and seminal vesicles that occurs during a seven day period after removal of the testes (orchiectomy [ORX]).

Rats are orchiectomized (ORX). Each rat is weighed, then anesthetized by isoflurane gas that is maintained to effect. A 1.5 cm anteroposterior incision is made in the scrotum. The right testicle is exteriorized. The spermatic artery and vas deferens are ligated with 4.0 silk 0.5cm proximal to the testicle. The testicle is freed by one cut of a small surgical scissors distal to the ligation site. The tissue stump is returned to the scrotum. The same is repeated for the left testicle. When both stumps are returned to the scrotum, the scrotum and overlying skin are sutured closed with 4.0 silk. For Sham-ORX, all procedures excepting ligation and scissors cutting are completed. The rats fully recover consciousness and full mobility within 10-15 minutes.

A dose of test compound is administered subcutaneously or orally to the rat immediately after the surgical incision is sutured. Treatment continues for an additional six consecutive days.

5    Necropsy and Endpoints:

The rat is first weighed, then anesthetized in a CO<sub>2</sub> chamber until near death. Approximately 5ml whole blood is obtained by cardiac puncture. The rat is then examined for certain signs of death and completeness of ORX. Next, the ventral portion of the prostate gland is located and blunt dissected free in a highly stylized  
10    fashion. The ventral prostate is blotted dry for 3-5 seconds and then weighed (VPW). Finally, the seminal vesicle is located and dissected free. The ventral seminal vesicle is blotted dry for 3-5 seconds and then weighed (SVWT).

Primary data for this assay are the weights of the ventral prostate and seminal vesicle. Secondary data include serum LH (luteinizing hormone) and FSH  
15    (follicle stimulating hormone), and possible serum markers of bone formation and virilization. Data are analyzed by ANOVA plus Fisher PLSD post-hoc test to identify intergroup differences. The extent to which test compounds inhibit ORX-induced loss of VPW and SVWT is assessed.

20    7. In Vivo Bone Formation Assay:

Female Sprague-Dawley rats aged 7-10 months are used in treatment mode to simulate adult human females. The rats have been ovariectomized (OVX) 75-180 days previously, to cause bone loss and simulate estrogen deficient, osteopenic adult human females. Pre-treatment with a low dose of a powerful anti-resorptive,  
25    alendronate (0.0028mpk SC, 2X/wk) is begun on Day 0. On Day 15, treatment with test compound is started. Test compound treatment occurs on Days 15-31 with necropsy on Day 32. The goal is to measure the extent to which androgen-like compounds increase the amount of bone formation, shown by increased fluorochrome labeling, at the periosteal surface.

30                    In a typical assay, nine groups of seven rats each are studied.

On Days 19 and 29 (fifth and fifteenth days of treatment), a single subcutaneous injection of calcein (8mg/kg) is given to each rat.

Necropsy and Endpoints:

The rat is first weighed, then anesthetized in a CO<sub>2</sub> chamber until near death. Approximately 5mL whole blood is obtained by cardiac puncture. The rat is then examined for certain signs of death and completeness of OVX. First, the uterus is located, blunt dissected free in a highly stylized fashion, blotted dry for 3-5 seconds and then weighed (UW). The uterus is placed in 10% neutral-buffered formalin. Next, the right leg is disarticulated at the hip. The femur and tibia are separated at the knee, substantially defleshed, and then placed in 70% ethanol.


A 1-cm segment of the central right femur, with the femoral proximal-distal midpoint at its center, is placed in a scintillation vial and dehydrated and defatted in graded alcohols and acetone, then introduced to solutions with increasing concentrations of methyl methacrylate. It is embedded in a mixture of 90% methyl methacrylate :10% dibutyl phthalate, that is allowed to polymerize over a 48-72hr period. The block is cracked and the plastic block is trimmed into a shape that conveniently fits the vice-like specimen holder of a Leica 1600 Saw Microtome, with the long axis of the bone prepared for cross-sectioning. Three cross-sections of 85 $\mu$ m thickness are prepared and mounted on glass slides. One section from each rat that approximates the midpoint of the bone is selected and blind-coded. The periosteal surface of each section is assessed for total periosteal surface, single fluorochrome label, double fluorochrome label, and interlabel distance.

Primary data for this assay are the percentage of periosteal surface bearing double label and the mineral apposition rate (interlabel distance( $\mu$ m)/10d), semi-independent markers of bone formation. Secondary data include uterus weight and histologic features. Tertiary endpoints may include serum markers of bone formation and virilization. Data are analyzed by ANOVA plus Fisher PLSD post-hoc test to identify intergroup differences. The extent to which test compounds increase bone formation endpoint are assessed.

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it is understood that the practice of the invention encompasses all of the usual variations, adoptions, or modifications, as being within the scope of the following claims and their equivalents.

## WHAT IS CLAIMED IS:

1. A compound of structural formula I:



5

or an enantiomer or a pharmaceutically acceptable salt thereof; wherein

n is 0, 1 or 2;

a-b represents CH=CH or CH<sub>2</sub>CH<sub>2</sub>;

10 "c" represents a double bond;

R<sup>1</sup> is selected from

hydrogen,

hydroxy,

C<sub>1-4</sub> alkoxy,

15

hydroxymethyl, and

C<sub>1-3</sub> alkyl, unsubstituted or substituted with one to seven fluorine atoms;

R<sup>4</sup> is selected from

halogen,

C<sub>1-6</sub> alkyl, wherein alkyl is unsubstituted or substituted with one to three

20

substituents independently selected from cyano, carboxy, halogen,

hydroxy, oxo, C<sub>1-4</sub> alkoxy, and C<sub>1-4</sub> alkylthio,

C<sub>2-6</sub> alkenyl,

C<sub>2-6</sub> alkynyl,

(CH<sub>2</sub>)<sub>n</sub>-phenyl, and

(CH<sub>2</sub>)<sub>n</sub>-naphthyl;

or R<sup>4</sup> together with the carbon atom to which it is attached form a carbonyl or cyclopropyl group, with the proviso that when R<sup>4</sup> together with the carbon atom to which it is attached form a carbonyl or cyclopropyl group, then "c" represents a single bond;

or R<sup>1</sup> and R<sup>4</sup> together with the atoms to which they are attached form a 5- or 6-membered ring system optionally containing an additional heteroatom selected from O, S, and NC<sub>1-4</sub> alkyl;

10 R<sup>2</sup> is hydrogen or C<sub>1-4</sub> alkyl;

R<sup>3</sup> is selected from

C<sub>1-4</sub> alkyl, wherein said C<sub>1-4</sub> alkyl is optionally substituted with one or more substituents independently selected from halogen, hydroxy, C<sub>1-4</sub> alkoxy, and C<sub>1-4</sub> alkylamino, and

15 (CH<sub>2</sub>)<sub>n</sub>-aryl, wherein said aryl is selected from:

(3) phenyl,

(2) naphthyl,

(3) benzimidazolyl,

(4) benzofuranyl,

20 (5) benzothiophenyl,

(6) benzoxazolyl,

(7) benzothiazolyl,

(8) benzodihydrofuranyl,

(9) 1,3-benzodioxolyl,

25 (10) 2,3-dihydro-1,4-benzodioxinyl,

(11) indolyl,

(12) quinolyl,

(13) isoquinolyl,

(14) furanyl,

- (15) thienyl,  
 (16) imidazolyl,  
 (17) oxazolyl,  
 (18) thiazolyl,  
 5 (19) isoxazolyl,  
 (20) isothiazolyl,  
 (21) pyrazolyl,  
 (22) pyrrolyl,  
 (23) pyridyl,  
 10 (24) pyrimidyl,  
 (25) pyrazinyl,  
 (26) thiadiazolyl,  
 (27) oxadiazolyl,  
 (28) triazolyl,  
 15 (29) imidizopyridinyl,  
 (30) tetrazolyl, and  
 (31) indanyl;

wherein the aryl group as defined in items (1) to (31) is unsubstituted or substituted  
 with one to three groups independently selected from halogen, phenyl,  
 20 C<sub>1-8</sub> alkyl, C<sub>3-8</sub> cycloalkyl, C<sub>3-8</sub> cycloheteroalkyl, phenyl-C<sub>1-6</sub> alkyl,  
 amino-C<sub>0-6</sub> alkyl, C<sub>1-6</sub> alkylamino-C<sub>0-6</sub> alkyl,  
 (C<sub>1-6</sub> alkyl)<sub>2</sub>amino-C<sub>0-6</sub> alkyl, phenyl-C<sub>0-6</sub> alkylamino-C<sub>0-6</sub> alkyl, (phenyl-  
 C<sub>0-6</sub> alkyl)<sub>2</sub>amino-C<sub>0-6</sub> alkyl, C<sub>1-6</sub> alkylthio, phenyl-C<sub>0-6</sub> alkylthio,  
 C<sub>1-6</sub> alkylsulfinyl, phenyl-C<sub>0-6</sub> alkylsulfinyl, C<sub>1-6</sub> alkylsulfonyl,  
 25 phenyl-C<sub>0-6</sub> alkylsulfonyl, C<sub>1-6</sub> alkoxy-C<sub>0-6</sub> alkyl,  
 phenyl-C<sub>0-6</sub> alkoxy-C<sub>0-6</sub> alkyl, hydroxycarbonyl-C<sub>0-6</sub> alkyl,  
 C<sub>1-6</sub> alkoxycarbonyl-C<sub>0-6</sub> alkyl, phenyl-C<sub>0-6</sub> alkoxycarbonyl-C<sub>0-6</sub> alkyl,  
 hydroxycarbonyl-C<sub>1-6</sub> alkyloxy, hydroxy-C<sub>0-6</sub> alkyl, cyano, nitro,  
 perfluoro-C<sub>1-4</sub> alkyl, perfluoro-C<sub>1-4</sub> alkoxy, C<sub>1-6</sub> alkylcarbonyloxy,

- phenyl-C<sub>0-6</sub> alkylcarbonyloxy, C<sub>1-6</sub> alkylcarbonylamino,  
 phenyl-C<sub>0-6</sub> alkylcarbonylamino, C<sub>1-6</sub> alkylsulfonylamino,  
 phenyl-C<sub>0-6</sub> alkylsulfonylamino, C<sub>1-6</sub> alkoxy carbonylamino,  
 phenyl-C<sub>0-6</sub> alkoxy carbonylamino, C<sub>1-6</sub> alkylaminocarbonylamino,  
 5 phenyl-C<sub>0-6</sub> alkylaminocarbonylamino, (C<sub>1-6</sub> alkyl)<sub>2</sub> aminocarbonylamino,  
 (phenyl-C<sub>0-6</sub> alkyl)<sub>2</sub> aminocarbonylamino, (C<sub>1-6</sub> alkyl)<sub>2</sub> aminocarbonyloxy,  
 and (phenyl-C<sub>0-6</sub> alkyl)<sub>2</sub> aminocarbonyloxy;  
 the alkyl group is unsubstituted or substituted with one to three substituents  
 independently selected from halogen, hydroxy, and C<sub>1-4</sub> alkoxy;  
 10 and wherein any methylene (CH<sub>2</sub>) carbon atom in (CH<sub>2</sub>)<sub>n</sub> is unsubstituted or  
 substituted with one to two groups independently selected from halogen,  
 hydroxy, and C<sub>1-4</sub> alkyl; or two substituents when on the same methylene  
 (CH<sub>2</sub>) group are taken together with the carbon atom to which they are  
 attached to form a cyclopropyl group;  
 15 or R<sup>2</sup> and R<sup>3</sup> together form a 5- or 6-membered saturated ring fused with a 5- or 6-  
 membered aromatic ring system having 0, 1, or 2 heteroatoms selected from  
 N, O, and S.

2. A compound of Claim 1, wherein:

- 20 n is 0, 1 or 2;  
 a-b represents CH=CH or CH<sub>2</sub>CH<sub>2</sub>;  
 "c" represents a double bond;  
 R<sup>1</sup> is selected from  
 hydrogen,  
 25 hydroxy,  
 C<sub>1-4</sub> alkoxy,  
 hydroxymethyl, and  
 C<sub>1-3</sub> alkyl, unsubstituted or substituted with one to seven fluorine atoms;  
 R<sup>4</sup> is selected from

halogen,

C<sub>1-6</sub> alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from cyano, carboxy, halogen, hydroxy, oxo, C<sub>1-4</sub> alkoxy, and C<sub>1-4</sub> alkylthio,

5 C<sub>2-6</sub> alkenyl,

C<sub>2-6</sub> alkynyl,

(CH<sub>2</sub>)<sub>n</sub>-phenyl, and

(CH<sub>2</sub>)<sub>n</sub>-naphthyl;

or R<sup>4</sup> together with the carbon atom to which it is attached form a carbonyl or

10 cyclopropyl group, with the proviso that when R<sup>4</sup> together with the carbon atom to which it is attached form a carbonyl or cyclopropyl group, then "c" represents a single bond;

or R<sup>1</sup> and R<sup>4</sup> together with the atoms to which they are attached form a 5- or 6-membered ring system optionally containing an additional heteroatom selected  
15 from O, S, and NC<sub>1-4</sub> alkyl;

R<sup>2</sup> is hydrogen or C<sub>1-4</sub> alkyl;

R<sup>3</sup> is selected from

C<sub>1-4</sub> alkyl, wherein said C<sub>1-4</sub> alkyl is optionally substituted with one or more substituents independently selected from halogen, hydroxy, C<sub>1-4</sub>  
20 alkoxy, and C<sub>1-4</sub> alkylamino, and

(CH<sub>2</sub>)<sub>n</sub>-aryl, wherein said aryl is selected from:

(4) phenyl,

(2) naphthyl,

(3) benzimidazolyl,

25 (4) benzofuranyl,

(5) benzothiophenyl,

(6) benzoxazolyl,

(7) benzothiazolyl,

(8) benzodihydrofuranyl,

- (9) 1,3-benzodioxolyl,  
 (10) 2,3-dihydro-1,4-benzodioxinyl,  
 (11) indolyl,  
 (12) quinolyl,  
 5 (13) isoquinolyl,  
 (14) furanyl,  
 (15) thienyl,  
 (16) imidazolyl,  
 (17) oxazolyl,  
 10 (18) thiazolyl,  
 (19) isoxazolyl,  
 (20) isothiazolyl,  
 (21) pyrazolyl,  
 (22) pyrrolyl,  
 15 (23) pyridyl,  
 (24) pyrimidyl,  
 (25) pyrazinyl,  
 (26) thiadiazolyl,  
 (27) oxadiazolyl,  
 20 (28) triazolyl,  
 (29) tetrazolyl, and  
 (30) indanyl;

wherein the aryl group as defined in items (1) to (31) is unsubstituted or substituted  
 with one to three groups independently selected from halogen, phenyl,  
 25 C<sub>1-8</sub> alkyl, C<sub>3-8</sub> cycloalkyl, C<sub>3-8</sub> cycloheteroalkyl, phenyl-C<sub>1-6</sub> alkyl,  
 amino-C<sub>0-6</sub> alkyl, C<sub>1-6</sub> alkylamino-C<sub>0-6</sub> alkyl,  
 (C<sub>1-6</sub> alkyl)<sub>2</sub>amino-C<sub>0-6</sub> alkyl, phenyl-C<sub>0-6</sub> alkylamino-C<sub>0-6</sub> alkyl, (phenyl-  
 C<sub>0-6</sub> alkyl)<sub>2</sub>amino-C<sub>0-6</sub> alkyl, C<sub>1-6</sub> alkylthio, phenyl-C<sub>0-6</sub> alkylthio,  
 C<sub>1-6</sub> alkylsulfinyl, phenyl-C<sub>0-6</sub> alkylsulfinyl, C<sub>1-6</sub> alkylsulfonyl,  
 30 phenyl-C<sub>0-6</sub> alkylsulfonyl, C<sub>1-6</sub> alkoxy-C<sub>0-6</sub> alkyl,

phenyl-C<sub>0-6</sub> alkoxy-C<sub>0-6</sub> alkyl, hydroxycarbonyl-C<sub>0-6</sub> alkyl,  
C<sub>1-6</sub> alkoxycarbonyl-C<sub>0-6</sub> alkyl, phenyl-C<sub>0-6</sub> alkoxycarbonyl-C<sub>0-6</sub> alkyl,  
hydroxycarbonyl-C<sub>1-6</sub> alkyloxy, hydroxy-C<sub>0-6</sub> alkyl, cyano, nitro,  
perfluoro-C<sub>1-4</sub> alkyl, perfluoro-C<sub>1-4</sub> alkoxy, C<sub>1-6</sub> alkylcarbonyloxy,  
5 phenyl-C<sub>0-6</sub> alkylcarbonyloxy, C<sub>1-6</sub> alkylcarbonylamino,  
phenyl-C<sub>0-6</sub> alkylcarbonylamino, C<sub>1-6</sub> alkylsulfonylamino,  
phenyl-C<sub>0-6</sub> alkylsulfonylamino, C<sub>1-6</sub> alkoxycarbonylamino,  
phenyl-C<sub>0-6</sub> alkoxycarbonylamino, C<sub>1-6</sub> alkylaminocarbonylamino,  
phenyl-C<sub>0-6</sub> alkylaminocarbonylamino, (C<sub>1-6</sub> alkyl)<sub>2</sub> aminocarbonylamino,  
10 (phenyl-C<sub>0-6</sub> alkyl)<sub>2</sub> aminocarbonylamino, (C<sub>1-6</sub> alkyl)<sub>2</sub> aminocarbonyloxy,  
and (phenyl-C<sub>0-6</sub> alkyl)<sub>2</sub> aminocarbonyloxy;

the alkyl group is unsubstituted or substituted with one to three substituents  
independently selected from halogen, hydroxy, and C<sub>1-4</sub> alkoxy;

and wherein any methylene (CH<sub>2</sub>) carbon atom in (CH<sub>2</sub>)<sub>n</sub> is unsubstituted or  
15 substituted with one to two groups independently selected from halogen,  
hydroxy, and C<sub>1-4</sub> alkyl; or two substituents when on the same methylene  
(CH<sub>2</sub>) group are taken together with the carbon atom to which they are  
attached to form a cyclopropyl group;

or R<sup>2</sup> and R<sup>3</sup> together form a 5- or 6-membered saturated ring fused with a 5- or 6-  
20 membered aromatic ring system having 0, 1, or 2 heteroatoms selected from  
N, O, and S;

or an enantiomer or a pharmaceutically acceptable salt thereof.

3. A compound of Claim 1, wherein R<sup>1</sup> is hydrogen or C<sub>1-3</sub> alkyl,  
25 unsubstituted or substituted with one to seven fluorine atoms.

4. A compound of Claim 3, wherein R<sup>1</sup> is hydrogen or methyl.

5. A compound of Claim 1, wherein a-b represents  $\text{CH}_2\text{CH}_2$ .
6. A compound of Claim 1, wherein a-b represents  $\text{CH}=\text{CH}$ .
- 5 7. A compound of Claim 4, wherein  $\text{R}^4$  is selected from  $\text{C}_{1-4}$   
alkyl.
8. A compound of Claim 7, wherein  $\text{R}^4$  is methyl.
9. A compound of Claim 8, wherein  $\text{R}^1$  is methyl.
- 10 10. A compound of Claim 4, wherein  $\text{R}^4$  together with the carbon  
atom to which it is attached form a carbonyl or cyclopropyl group.
11. A compound of Claim 10, wherein  $\text{R}^4$  together with the carbon  
15 atom to which it is attached is a cyclopropyl group.
12. A compound of Claim 11, wherein  $\text{R}^1$  is hydrogen or methyl.
13. A compound of Claim 1, wherein  $\text{R}^1$  and  $\text{R}^4$  together with the  
20 atoms to which they are attached form a 5- or 6-membered ring system optionally  
containing an additional heteroatom selected from O, S, and  $\text{NC}_{1-4}$  alkyl.
14. A compound of Claim 1, wherein  $\text{R}^2$  is hydrogen.
- 25 15. A compound of Claim 14, wherein  $\text{R}^3$  is  $(\text{CH}_2)_n$ -aryl.
16. A compound of Claim 15, wherein  $\text{R}^3$  is  $(\text{CH}_2)_n$ -aryl and n is  
selected from 0 and 1, and aryl is selected from: phenyl, imidazopyridinyl,  
benzimidazolyl, benzothiophenyl, indolyl, quinolyl, isoquinolyl, thienyl, imidazolyl,

thiazolyl, pyrazolyl, pyrrolyl, pyridyl, pyrimidyl, pyrazinyl, thiadiazolyl, triazolyl, and indanyl.

17. A compound of Claim 16, wherein aryl is selected from:  
 5 phenyl, benzimidazolyl, benzothiophenyl, indolyl, pyrazolyl, pyrrolyl, pyridyl, pyrimidyl, pyrazinyl, thiadiazolyl, imidazopyridinyl, and triazolyl.

18. A compound of Claim 17, wherein R<sup>1</sup> is methyl.

19. A compound of Claim 18, wherein R<sup>4</sup> is selected from C<sub>1-4</sub> alkyl, wherein alkyl is unsubstituted or substituted with one to three substituents independently selected from cyano, carboxy, halogen, hydroxy, oxo, C<sub>1-4</sub> alkoxy, and C<sub>1-4</sub> alkylthio, (CH<sub>2</sub>)<sub>n</sub>-phenyl, and halogen.

20. A compound of Claim 19, wherein R<sup>4</sup> is a methyl group.

21. A compound of Claim 17, wherein R<sup>4</sup> together with the carbon atom to which it is attached form a cyclopropyl group, and R<sup>1</sup> is selected from hydrogen and methyl.

20

22. A compound of Claim 1 selected from:  
 N-phenyl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(2-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
 carboxamide,  
 25 N-(3-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
 carboxamide,  
 N-(4-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
 carboxamide,  
 N-(2-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
 30 carboxamide,

- N-(3-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
5 N-(2-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(3-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(4-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
10 carboxamide,  
N-(2,2,2-trifluoroethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(2-fluorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
15 N-(3-fluorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(4-fluorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
20 carboxamide,  
N-(2,3-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-  
5-en-17 $\beta$ -carboxamide,  
N-(2,5-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-  
5-en-17 $\beta$ -carboxamide,  
25 N-(2-chloro-6-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-  
en-17 $\beta$ -carboxamide,  
N-(2-chloro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-  
en-17 $\beta$ -carboxamide,  
N-(4-chloro-2-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-  
30 en-17 $\beta$ -carboxamide,

- N-(2,4-dichloro-5-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-[4-(morpholin-4-yl)phenyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
5 N-(3-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(4-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
10 carboxamide,  
N-[2-(piperidin-1-yl)phenyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(5-fluoro-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
15 N-(2-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(4-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(5-methyl-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
20 carboxamide,  
N-(4-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
25 N-(3-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(3-cyanophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
30 N-(pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

- N-(pyridin-3-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(2-chloro-4-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
5 N-(2,6-dichlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(2,6-difluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(2,3,4,6-tetrafluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
10 carboxamide,  
N-(2,3-dichlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(2,5-dichlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
15 N-(thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(3-chloro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
20 N-(3-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(3-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(4-acetyl-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
25 carboxamide,  
N-(4-methyl-thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(4-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,

- N-(2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2,5-difluoro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-  
17 $\beta$ -carboxamide,
- 5 N-(2,4,5,6-tetrafluoro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,6-dichloro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-(3,4-dicyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,4-dichloro-6-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-chloro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-(3-fluoro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3-trifluoromethyl-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-chloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 20 N-(3,5-dichloro-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-chloro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-(2,5-dichloro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3-chloro-5-trifluoromethyl-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,6-dichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 30

- 3-[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino}-4-chlorobenzoic acid,  
 N-(4-chlorobenzothiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 5 N-(benzothiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(4,5-dimethylthiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(2,3,5-trichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 10 N-(2,6-dichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(5-ethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 15 N-(4-methyl-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(2-chloro-6-trifluoromethyl-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 20 N-(2,6-dichloro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(2,6-dichloro-6'-trifluoro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 25 N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(4-methyl-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(pyrazin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

- N-(6-chloro-pyrazin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
5 N-(2-fluoro-6-chlorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-  
17 $\beta$ -carboxamide,  
N-((R)-phenylmethylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-((S)-phenylmethylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
10 carboxamide,  
N-(2,6-dichlorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(3,4-dichlorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
15 N-[(1S,2S)-2-hydroxycyclohexyl] -4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-  
17 $\beta$ -carboxamide,  
N-(3H-imidazo[4,5-b]pyridin-2-ylmethyl) )-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -  
androst-5-en-17 $\beta$ -carboxamide,  
N-[(1R,2S)-2-hydroxycyclohexyl] -4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-  
20 17 $\beta$ -carboxamide,  
N-(1,3-thiazol-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(1H-indol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
25 N-[2-(1H-benzimidazol-2-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-  
17 $\beta$ -carboxamide,  
N-[2-(6-chloro-1H-benzimidazol-2-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -  
androst-5-en-17 $\beta$ -carboxamide,

- tert-butyl 3-({[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino}methyl)-1H-indole-1-carboxylate,
- 5 tert-butyl 3-(2-{{[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino}ethyl)-1H-indole-1-carboxylate,
- N-(benzimidazol-2-ylmethylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-[2-(1-methyl-1H-benzimidazol-2-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(imidazo[1,2-a]pyrimidin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(1H-indol-5-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-[(6-chloro-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(imidazo[1,2-a]pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-imidazo[1,2-a]pyridin-2-ylethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 20 N-[2-(2-phenyl-1H-indol-3-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(1H-indol-7-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-(1H-indol-4-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(imidazo[1,2-a]pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[(1-phenyl-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 30

- N-(pyrazolo[1,5-a]pyridin-7-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(pyrazolo[1,5-a]pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
5 N-[(1-methyl-2,3-dihydro-1H-indol-5-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-piperidin-4-yl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-piperidin-3-yl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2-ethylpyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
10 carboxamide,  
N-[(1-methyl-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(5-t-Butyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
15 N-(5-trifluoromethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2-methoxypyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2-ethoxypyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
20 carboxamide,  
N-(pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(4-trifluoromethylpyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(4,6-dimethylpyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
25 carboxamide,  
N-(6-methylpyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(7H-purin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

- N-(5-cyclopropyl-1*H*-pyrazol-3-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(6-methoxybenzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-(2,3-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2,6-dichlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-fluoro-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 10 N-(2-fluoro-6-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-trifluoromethyl-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-fluoro-4-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-trifluoromethylphenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 15 N-(2-fluoro-5-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-phenyl-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(5-ethyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(5-methyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 20 N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(pyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-chloropyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 25 N-(2-methylpyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-chloropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(4-fluoropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(4-chloropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(4-methylpyridin-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 30 N-(4-chloropyridin-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

- N-(benzimidazol-2-ylmethyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
N-((N-methylbenzimidazol-2-yl)methyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
5 N-(3H-imidazo[4,5-b]pyridin-2-ylmethyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
N-(pyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2,3-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
10 N-(2-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2,6-dichlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
N-(2-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-fluoro-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -  
15 carboxamide,  
N-(2,6-difluorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
N-(2-trifluoromethyl-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -  
androst-17 $\beta$ -carboxamide,  
20 N-(2-fluoro-4-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
N-(2-trifluoromethylphenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
N-(2-fluoro-5-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -  
25 carboxamide,  
N-phenyl-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2,6-dichloro-4-cyanophenyl) -6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-  
17 $\beta$ -carboxamide,  
N-(1-methyl-6-trifluoromethyl-1H-benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-  
30 oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide.

N-(2-methyl-3-chloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
enantiomers, and pharmaceutically acceptable salts thereof.

- 5                    23.     A compound of Claim 22 selected from:  
N-(3H-imidazo[4,5-b]pyridin-2-ylmethyl) )-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(imidazo[1,2-a]pyrimidin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
10 N-(imidazo[1,2-a]pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2-imidazo[1,2-a]pyridin-2-ylethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(3H-imidazo[4,5-b]pyridin-2-ylmethyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
15 carboxamide,  
enantiomers, and pharmaceutically acceptable salts thereof..

24.     A compound of Claim 23 selected from:  
N-(3H-imidazo[4,5-b]pyridin-2-ylmethyl) )-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
20 N-(imidazo[1,2-a]pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(3H-imidazo[4,5-b]pyridin-2-ylmethyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
25 enantiomers, and pharmaceutically acceptable salts thereof.

25.     A compound of Claim 22 selected from:  
N-phenyl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
30 carboxamide,

- N-(3-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(4-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- 5 N-(2-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(3-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
10 carboxamide,
- N-(2-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(3-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- 15 N-(4-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2,2,2-trifluoroethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2-fluorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
20 carboxamide,
- N-(3-fluorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(4-fluorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- 25 N-(pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2,3-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-  
5-en-17 $\beta$ -carboxamide,
- N-(2,5-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-  
30 5-en-17 $\beta$ -carboxamide,

- N-(2-chloro-6-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-chloro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-(4-chloro-2-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,4-dichloro-5-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[4-(morpholin-4-yl)phenyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-(3-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(4-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-(2-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[2-(piperidin-1-yl)phenyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(5-fluoro-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 20 N-(2-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(4-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-(5-methyl-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(4-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 30

- N-(3-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(3-cyanophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- 5 N-(pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(pyridin-3-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2-chloro-4-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
10 carboxamide,
- N-(2,6-dichlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2,6-difluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- 15 N-(2,3,4,6-tetrafluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2,3-dichlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2,5-dichlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
20 carboxamide,
- N-(thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(3-chloro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
25 carboxamide,
- N-(3-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(3-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,

- N-(4-acetyl-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(4-methyl-thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-(4-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,5-difluoro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-(2,4,5,6-tetrafluoro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,6-dichloro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-(3,4-dicyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,4-dichloro-6-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-chloro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 20 N-(3-fluoro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3-trifluoromethyl-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-(2-chloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3,5-dichloro-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-chloro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 30

- N-(2,5-dichloro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3-chloro-5-trifluoromethyl-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-(2,6-dichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 3-{[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino}-4-chlorobenzoic acid,
- N-(4-chlorobenzothiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-(benzothiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(4,5-dimethylthiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-(2,3,5-trichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,6-dichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(5-ethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 20 N-(4-methyl-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-(2-chloro-6-trifluoromethyl-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,6-dichloro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,6-dichloro-6'-trifluoro-pyridin-3-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 30

- N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(4-methyl-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
5 N-(pyrazin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(6-chloro-pyrazin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
10 N-(2-fluoro-6-chlorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-((R)-phenylmethylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-((S)-phenylmethylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
15 N-(2,6-dichlorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(3,4-dichlorophenylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
20 N-[(1S,2S)-2-hydroxycyclohexyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-[(1R,2S)-2-hydroxycyclohexyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(1,3-thiazol-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
25 N-(1H-indol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-[2-(1H-benzimidazol-2-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

- N-[2-(6-chloro-1H-benzimidazol-2-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 tert-butyl 3-({[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino}methyl)-1H-  
 5 indole-1-carboxylate,  
 tert-butyl 3-(2-({[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino}ethyl)-1H-indole-1-carboxylate,  
 N-(benzimidazol-2-ylmethylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 10 N-[2-(1-methyl-1H-benzimidazol-2-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(1H-indol-5-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 15 N-[(6-chloro-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-[2-(2-phenyl-1H-indol-3-yl)ethyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(1H-indol-7-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
 20 carboxamide,  
 N-(1H-indol-4-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-[(1-phenyl-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 25 N-(pyrazolo[1,5-a]pyridin-7-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(pyrazolo[1,5-a]pyridin-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-[(1-methyl-2,3-dihydro-1H-indol-5-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -  
 30 androst-5-en-17 $\beta$ -carboxamide,

- N-piperidin-4-yl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-piperidin-3-yl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2-ethylpyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
5 N-[(1-methyl-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -  
androst-5-en-17 $\beta$ -carboxamide,  
N-(5-t-Butyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-  
17 $\beta$ -carboxamide,  
N-(5-trifluoromethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -  
10 androst-5-en-17 $\beta$ -carboxamide,  
N-(2-methoxypyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(2-ethoxypyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
15 N-(pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(4-trifluoromethylpyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-  
17 $\beta$ -carboxamide,  
N-(4,6-dimethylpyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
20 N-(6-methylpyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(7H-purin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(5-cyclopropyl-1H-pyrazol-3-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -  
25 androst-5-en-17 $\beta$ -carboxamide,  
N-(6-methoxybenzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-  
5-en-17 $\beta$ -carboxamide,  
N-(1-methyl-6-trifluoromethyl-1H-benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-  
oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
30 N-(2,3-Difluorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

- N-(2-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(2,6-dichlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(2-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(2-fluoro-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 5 N-(2-fluoro-6-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(2-trifluoromethyl-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
 carboxamide,  
 N-(2-fluoro-4-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(2-trifluoromethylphenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 10 N-(2-fluoro-5-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-phenyl-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(5-ethyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
 carboxamide,  
 N-(5-methyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
 15 carboxamide,  
 N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
 carboxamide,  
 N-(pyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(2-chloropyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 20 N-(2-methylpyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(2-chloropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(4-fluoropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(4-chloropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(4-methylpyridin-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 25 N-(4-chloropyridin-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
 N-(benzimidazol-2-ylmethyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
 carboxamide,  
 N-((N-methylbenzimidazol-2-yl)methyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
 carboxamide,  
 30 N-(pyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

- N-(2,3-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2,6-dichlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 5 N-(2-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2-fluoro-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2,6-difluorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 10 N-(2-trifluoromethyl-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2-fluoro-4-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 15 N-(2-trifluoromethylphenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2-fluoro-5-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-phenyl-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- 20 N-(2,6-dichloro-4-cyanophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,
- N-(2-methyl-3-chloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- enantiomers, and pharmaceutically acceptable salts thereof.

25

26. A compound of Claim 25 selected from:

- N-phenyl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

- N-(3-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(3-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- 5 N-(2,3-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-  
5-en-17 $\beta$ -carboxamide,
- N-(2,5-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-  
5-en-17 $\beta$ -carboxamide,
- 10 N-(3-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(3-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- 15 N-(3-cyanophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2,6-difluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2,3,4,6-tetrafluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
20 carboxamide,
- N-(thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(4-methyl-thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
25 carboxamide,
- N-(4-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,
- N-(2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,

- N-(2-chloro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3-fluoro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 5 N-(2-chloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3,5-dichloro-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(4,5-dimethylthiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-(2,3,5-trichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2,6-dichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-(5-ethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(4-methyl-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 20 N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(6-chloro-pyrazin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-(benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[(1S,2S)-2-hydroxycyclohexyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-[(1R,2S)-2-hydroxycyclohexyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 30

- tert-butyl 3-({[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino)methyl)-1H-indole-1-carboxylate,
- 5 N-[(6-chloro-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(1-methyl-6-trifluoromethyl-1H-benzimidazol-2-yl)methyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-ethylpyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 10 N-[(1-methyl-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(5-t-Butyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(5-trifluoromethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 15 N-(2-methoxypyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(2-methyl-3-chloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 20 enantiomers, and pharmaceutically acceptable salts thereof.

27. A compound of Claim 26 selected from:

- N-phenyl-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3-chlorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- 25 N-(3-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,
- N-(3-methoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,

- N-(2,3-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2,5-difluoro-4-trifluoromethylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
5 N-(3-trifluoromethoxyphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2-fluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(3-methylphenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
10 N-(3-cyanophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2,6-difluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
15 N-(2,3,4,6-tetrafluorophenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(3-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(4-chloro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
20 N-(2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(2-chloro-4-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
25 N-(3-fluoro-2-cyano-phenyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-(benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
N-[(1S,2S)-2-hydroxycyclohexyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
30

- N-[(1R,2S)-2-hydroxycyclohexyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 tert-butyl 3-({[(1,4a,6a,11-tetramethyl-2-oxo-2,3,4,4a,4b,5,6,6a,7,8,9,9a,9b,10-tetradecahydro-1H-indeno[5,4-f]quinolin-7-yl)carbonyl]amino }methyl)-1H-indole-1-carboxylate,  
 5 N-[(6-chloro-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(1-methyl-6-trifluoromethyl-1H-benzimidazol-2-ylmethyl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 10 N-[(1-methyl-1H-benzimidazol-2-yl)methyl]-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 enantiomers, and pharmaceutically acceptable salts thereof.

28. A compound of Claim 26 selected from:

- 15 N-(thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(4-methyl-thiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(4,5-dimethylthiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 20 N-(5-ethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(5-t-Butyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(5-trifluoromethyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 25 N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 N-(5-methyl-1,3,4-thiadiazol-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -carboxamide,  
 30 enantiomers, and pharmaceutically acceptable salts thereof.

29. A compound of Claim 26 selected from:

- N-(2-chloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
5 N-(3,5-dichloro-pyridin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(2,3,5-trichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
10 N-(2,6-dichloro-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(4-methyl-pyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(6-chloro-pyrazin-2-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
15 N-(2-ethylpyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
N-(2-methoxypyridin-4-yl)-4-methyl-6-methyl-3-oxo-4-aza-5 $\alpha$ -androst-5-en-17 $\beta$ -  
carboxamide,  
enantiomers, and pharmaceutically acceptable salts thereof.

20

30. A compound of claim 25 selected from:

- N-(2,3-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2,6-dichlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
25 N-(2-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-fluoro-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-fluoro-6-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-trifluoromethyl-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
30 N-(2-fluoro-4-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

- N-(2-trifluoromethylphenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-fluoro-5-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-phenyl-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(5-ethyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
5 carboxamide,  
N-(5-methyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
10 N-(pyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-chloropyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-methylpyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-chloropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(4-fluoropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
15 N-(4-chloropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(4-methylpyridin-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(4-chloropyridin-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(benzimidazol-2-ylmethyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
20 N-((N-methylbenzimidazol-2-yl)methyl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
N-(pyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2,3-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
25 N-(2-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2,6-dichlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
N-(2-fluorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(2-fluoro-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -  
30 carboxamide,

N-(2,6-difluorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-(2-trifluoromethyl-6-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,

5 N-(2-fluoro-4-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-(2-trifluoromethylphenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,

10 N-(2-fluoro-5-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-phenyl-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-(2,6-dichloro-4-cyanophenyl) -6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -carboxamide,

enantiomers, and pharmaceutically acceptable salts thereof.

15

31. A compound of claim 30 selected from:

N-(5-ethyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

20 N-(5-methyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

enantiomers, and pharmaceutically acceptable salts thereof.

25

32. A compound of Claim 30 selected from:

N-(pyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-(2-chloropyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-(2-methylpyridin-4-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-(2-chloropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

30 N-(4-fluoropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,

N-(4-chloropyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(4-methylpyridin-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(4-chloropyridin-2-yl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(pyridin-3-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
5 enantiomers, and pharmaceutically acceptable salts thereof.

33. A compound of Claim 30 selected from:  
N-(2,3-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
N-(4-methylpyridin-2-yl) -6,6-Ethylene-3-oxo-4-aza-5 $\alpha$ -androst-17 $\beta$ -carboxamide,  
10 N-(2,3-Difluoro-phenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl-5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
N-(2-fluoro-5-chlorophenyl)-6,6-Ethylene-3-oxo-4-aza-4-methyl -5 $\alpha$ -androst-17 $\beta$ -  
carboxamide,  
enantiomers, and pharmaceutically acceptable salts thereof.

15  
34. A method for modulating a function mediated by the androgen  
receptor in a mammal in need of such modulation comprising administering a  
therapeutically effective amount of a compound of Claim 1 or an enantiomer or a  
pharmaceutically acceptable salt thereof.

20  
35. A method of activating the function of the androgen receptor in  
a mammal in need of such activation comprising administering a therapeutically  
effective amount of a compound of Claim 1 or an enantiomer or a pharmaceutically  
acceptable salt thereof.

25  
36. A method of Claim 34, wherein said function mediated by the  
androgen receptor is activated in bone and/or muscle tissue and blocked in the  
prostate or the uterus.

37. A method of treating a condition in a mammal which is caused by androgen deficiency or which can be ameliorated by androgen replacement, which condition is selected from osteoporosis, osteopenia, glucocorticoid-induced osteoporosis, periodontal disease, bone fracture, bone damage following bone reconstructive surgery, sarcopenia, frailty, aging skin, male hypogonadism, postmenopausal symptoms in women, atherosclerosis, hypercholesterolemia, hyperlipidemia, obesity, aplastic anemia and other hematopoietic disorders, inflammatory arthritis and joint repair, HIV-wasting, prostate cancer, cancer cachexia, muscular dystrophies, Alzheimer's disease, premature ovarian failure, and autoimmune disease, comprising administering to the mammal in need of such treatment, a therapeutically effective amount of a compound of formula I or an enantiomer or a pharmaceutically acceptable salt thereof.

38. A method according to Claim 37, wherein said condition is osteoporosis.

39. A method of treating osteoporosis in a mammal in need thereof, comprising administering a therapeutically effective amount of a compound according to Claim 1 or an enantiomer or a pharmaceutically acceptable salt thereof.

40. A method of Claim 39 further comprising the administration of an agent selected from:

- (a) an estrogen or an estrogen derivative, alone or in combination with a progestin or progestin derivative,
- (b) a bisphosphonate,
- (c) an antiestrogen or a selective estrogen receptor modulator,
- (d) an  $\alpha_v\beta_3$  integrin receptor antagonist,
- (e) a cathepsin K inhibitor,
- (f) an HMG-CoA reductase inhibitor,
- (g) an osteoclast vacuolar ATPase inhibitor,

- (h) an antagonist of VEGF binding to osteoclast receptors,
- (i) an activator of peroxisome proliferator-activated receptor  $\gamma$ ,
- (j) calcitonin,
- (k) a calcium receptor antagonist,
- 5 (l) parathyroid hormone or analog thereof,
- (m) a growth hormone secretagogue,
- (n) human growth hormone,
- (o) insulin-like growth factor,
- (p) a p38 protein kinase inhibitor,
- 10 (q) bone morphogenetic protein,
- (r) an inhibitor of BMP antagonism,
- (s) a prostaglandin derivative,
- (t) vitamin D or vitamin D derivative,
- (u) vitamin K or vitamin K derivative,
- 15 (v) ipriflavone,
- (w) fluoride salts,
- (x) dietary calcium supplement, and
- (y) osteoprotegerin.

20

41. A method according to Claim 40, wherein said agent is selected from:

- (a) the estrogen or estrogen derivative, alone or in combination with a progestin or progestin derivative, is selected from conjugated estrogen, equine estrogen, 25 17 $\beta$ -estradiol, estrone, 17 $\beta$ -ethynyl estradiol, 17 $\beta$ -ethynyl estradiol with at least one compound selected from norethindrone and medroxyprogesterone acetate;
- (b) the bisphosphonate is selected from alendronate, clodronate, etidronate, ibandronate, incadronate, minodronate, neridronate, olpadronate, pamidronate, 30 piridronate, risedronate, tiludronate, and zoledronate;

- (c) the antiestrogen or selective estrogen receptor modulator is selected from raloxifene, clomiphene, zuclomiphene, enclomiphene, nafoxidene, CI-680, CI-628, CN-55,945-27, Mer-25, U-11,555A, U-100A, tamoxifen, lasofoxifene, toremifene, azorxifene, EM-800, EM-652, TSE 424, droloxifene, idoxifene, and levormeloxifene;
- (d) the HMG-CoA reductase inhibitor is selected from lovastatin, simvastatin, dihydroxy-open acid simvastatin, pravastatin, fluvastatin, atorvastatin, cerivastatin, rosuvastatin, pitavastatin, and nisvastatin;
- (e) calcitonin is salmon calcitonin administered as a nasal spray;
- 10 (f) bone morphogenetic protein is selected from BMP 2, BMP 3, BMP 5, BMP 6, BMP 7, TGF beta, and GDF5;
- (g) insulin-like growth factor is selected from IGF I and IGF II alone or in combination with IGF binding protein 3;
- (h) the prostaglandin derivative is selected from agonists of prostaglandin receptors EP1, EP2, EP4, FP, and IP;
- 15 (i) the fibroblast growth factor is selected from aFGF and bFGF;
- (j) parathyroid hormone (PTH) or PTH analog is selected from PTH subcutaneous injection, human PTH (1-84), human PTH (1-34), and other partial sequences, native or with substitutions;
- 20 (k) vitamin D or vitamin D derivative is selected from natural vitamin D, 25-OH-vitamin D3, 1 $\mu$ ,25(OH)<sub>2</sub> vitamin D3, 1 $\mu$ -OH-vitamin D3, 1 $\mu$ -OH-vitamin D2, dihydrotachysterol, 26,27-F6-1 $\mu$ ,25(OH)<sub>2</sub> vitamin D3, 19-nor-1 $\mu$ ,25(OH)<sub>2</sub> vitamin D3, 22-oxacalcitriol, calcipotriol, 1 $\mu$ ,25(OH)<sub>2</sub>-16-ene-23-yne-vitamin D3 (Ro 23-7553), EB1089, 20-epi-1 $\mu$ ,25(OH)<sub>2</sub> vitamin D3, KH1060, ED71, 1 $\mu$ ,24(S)-(OH)<sub>2</sub> vitamin D3, and 1 $\mu$ ,24(R)-(OH)<sub>2</sub> vitamin D3;
- 25 (l) the dietary calcium supplement is selected from calcium carbonate, calcium citrate, and natural calcium salts; and
- (m) the fluoride salts are selected from sodium fluoride and monosodium fluorophosphate (MFP);

and pharmaceutically acceptable salts or enantiomers thereof.

42. A method of Claim 41, wherein the bisphosphonate is alendronate monosodium trihydrate or alendronate monosodium monohydrate.

5

43. A method of Claim 40, wherein the agent is selected from:

- (a) an estrogen or an estrogen derivative, alone or in combination with a progestin or progestin derivative,
- (b) a bisphosphonate,
- 10 (c) an antiestrogen or a selective estrogen receptor modulator,
- (d) an  $\alpha v \beta 3$  integrin receptor antagonist,
- (e) a cathepsin K inhibitor,
- (f) an osteoclast vacuolar ATPase inhibitor,
- (g) calcitonin,
- 15 (h) parathyroid hormone or analog thereof,
- (i) and osteoprotegerin.

44. A pharmaceutical composition comprising a therapeutically effective amount of a compound of Claim 1 and a pharmaceutically acceptable carrier.

20

45. A composition of Claim 44 which further comprises an active ingredient selected from:

- (a) an estrogen or an estrogen derivative, alone or in combination with a progestin or progestin derivative;
- 25 (b) a bisphosphonate;
- (c) an antiestrogen or a selective estrogen receptor modulator,
- (d) an  $\alpha v \beta 3$  integrin receptor antagonist,
- (e) a cathepsin K inhibitor,
- (f) an HMG-CoA reductase inhibitor,
- 30 (g) an osteoclast vacuolar ATPase inhibitor,

- (h) an antagonist of VEGF binding to osteoclast receptors,  
(i) an activator of peroxisome proliferator-activated receptor  $\gamma$ ,  
(j) calcitonin,  
(k) a calcium receptor antagonist,  
5 (l) parathyroid hormone or analog thereof,  
(m) a growth hormone secretagogue,  
(n) human growth hormone,  
(o) insulin-like growth factor,  
(p) a p38 protein kinase inhibitor,  
10 (q) bone morphogenetic protein,  
(r) an inhibitor of BMP antagonism,  
(s) a prostaglandin derivative,  
(t) vitamin D or vitamin D derivative,  
(u) vitamin K or vitamin K derivative,  
15 (v) ipriflavone,  
(w) fluoride salts,  
(x) dietary calcium supplement, and  
(y) osteoprotegerin.

20 46. A composition of Claim 45, wherein said bisphosphonate is  
alendronate monosodium trihydrate or alendronate monosodium monohydrate.

47. A pharmaceutical composition made by combining a  
compound according to Claim 1 and a pharmaceutically acceptable carrier.

25

48. A process for making a pharmaceutical composition  
comprising combining a compound according to Claim 1 and a pharmaceutically  
acceptable carrier.

30